

DISS. ETH Nr. 18389

**The localization, protein interaction, and cell surface mobility  
of neuronal  $\alpha 7$  nicotinic acetylcholine receptors**

ABHANDLUNG

zur Erlangung des Titels

DOKTOR DER WISSENSCHAFTEN

der

ETH ZÜRICH

vorgelegt von

Thomas Bürli

Dipl. Natw. ETH

geboren am 24.02.1979

von

Klingnau AG

Angenommen auf Antrag von

Prof. Dr. Isabelle Mansuy, Referentin  
Prof. Dr. Ulrich Suter, Korreferent  
Dr. Christian Fuhrer, Korreferent  
Prof. Dr. Jean-Marc Fritschy, Korreferent

2009

## **Acknowledgements**

The work of this thesis was done in two parts. The first part was done during December 2004 to August 2006 in the Department of Neurochemistry of the Brain Research Institute at the Neuroscience Centre Zürich (University of Zürich and Swiss Federal Institute of Technology Zürich). The second part was done during August 2006 to May 2009 in the Department of Neuromorphology in the Institute of Pharmacology and Toxicology at the University of Zürich.

I thank Dr. Christian Fuhrer for supervising me in the first two years of my work. He supported me during this time in many aspects. He introduced me patiently into the field of neuroscience and he explained me the missing neuroscience background which I did not get in my education. During my work I profited from his trust and the freedom he gave me. He was always open and supportive when I walked in his office with another idea of an experiment. This attitude paired with his belief in success, especially in times when useful results were rare, enabled finally to succeed.

I thank Prof. Dr. Jean-Marc Fritschy for making it possible to continue my project in his group. Even though my work was not related to his field, he enabled me the best integration possible into his group. It was a pleasure to work in such a well equipped lab. Most of my findings would not exist without the high-end microscopes and the enormous palette of antibodies provided in his lab. Besides the material support, I would like to thank him for always having his door open for scientific and personal needs. His ability to ask always the right questions helped me to open my eyes and to see my project not as an isolated unit but as a piece embedded in a big puzzle, called neuroscience.

During the past four years, Dr. Kristin Baer was accompanying and supporting my project. I would like to thank her for always having time to discuss my work whenever it was necessary. Even though she is living in Swansea (Wales, UK) she was always well integrated in my project and we managed to exchange our ideas via Skype.

My work would not have been possible without the help of our technicians. Dubravka Göckeritz-Dujmovic, Corinne Sidler, Conny Schwerdel, and Ruth

Keist supported me with their excellent work and open minded attitude. I could always count on their big experience especially in primary cell culture and molecular cloning issues.

I would like to thank my lab office mate Caroline Petitjean for the excellent time we had. She was the one explaining and helping me whenever something was unclear how a microscope is built and how it works. She motivated me to discuss our work from a broader point of view, detached from facts and enriched with adventurous speculations.

Further, I thank Martina Baldinger. Together with Caroline they questioned my results frequently and their input was always very helpful to increase the quality of my work.

Dr. Shiva Tyagarajan was the advisor whenever I needed help in biochemical issues. I was always fascinated with how many helpful and inspiring ideas and experiments he came up just like that.

Aside from my lab I thank the whole pharmacological institute for the wonderful and inspiring time I had. The variety of interesting ongoing projects as well as the social bonding made it a pleasure to work in this institute.

Outside the institute, my thanks go to my friends who were a great support, especially in times when my work was not very productive. Special thanks go to my parents. They always had trust in me and they supported me in any imaginable way.

---

## Table of Contents

<b>Acknowledgements</b>	<b>2</b>
<b>Table of Contents</b>	<b>5</b>
<b>Table of Abbreviations</b>	<b>6</b>
<b>Summary</b>	<b>7</b>
<b>Zusammenfassung</b>	<b>9</b>
<b>1. Introduction</b>	<b>11</b>
1.1. Discovery of the acetylcholine receptor	11
1.2. Principles of neurotransmission	14
1.3. From static to dynamic view	22
1.4. Nicotinic acetylcholine receptors in the brain	26
1.5. Aim	34
<b>2. PICK1 interacts with <math>\alpha 7</math> neuronal nicotinic acetylcholine receptors and controls their clustering</b>	<b>37</b>
2.1. Abstract	38
2.2. Introduction	38
2.3. Results	40
2.4. Discussion	57
2.5. Experimental methods	64
2.6. Acknowledgements	71
<b>3. Efficient transfection of DNA or shRNA vectors into neurons using magnetofection</b>	<b>73</b>
3.1. Abstract	74
3.2. Introduction	74
3.3. Materials	76
3.4. Procedure	79
3.5. Anticipated results	87
3.6. Acknowledgements	97
<b>4. Single particle tracking of <math>\alpha 7</math> nicotinic acetylcholine receptors in hippocampal neurons reveals regulated perisynaptic confinement at glutamatergic and GABAergic perisynaptic sites</b>	<b>99</b>
4.1. Abstract	100
4.2. Introduction	101
4.3. Experimental procedures	103
4.4. Results	107
4.5. Discussion	118
4.6. Acknowledgements	123
<b>5. Discussion</b>	<b>125</b>
5.1. Summary of the results	125
5.2. In agreement or controversial?	126
5.3. What role does PICK1 play?	128
5.4. <i>In vivo</i> significance of the present results	130
5.5. Perspectives	134
<b>References</b>	<b>137</b>
<b>Curriculum Vitae</b>	<b>146</b>
<b>List of Publications</b>	<b>147</b>

## Table of Abbreviations

5-HT <sub>3</sub> R	5-hydroxytryptamine <sub>3</sub> or serotonin receptor	KO	knock out
α-BT	alpha bungarotoxin	LTP	long term potentiation
α7 nAChR	alpha7 nicotinic acetylcholine receptor	mAChR	muscarinic acetylcholine receptors
aa	amino acid	MEM	minimal essential medium
AA	Alanine-Alanine	MLA	methyllycaconitine citrate hydrate
ACh	acetylcholine	MSD	mean square displacement
AChE	acetylcholine esterase	nAChR	nicotinic acetylcholine receptor
AD	Alzheimer's disease	NGS	normal goat serum
AF	Alexa Fluor® dye	NMDAR	N-methyl D-aspartate receptor
AMPA	α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor	NMJ	neuromuscular junction
AP	action potential	PBS	phosphate buffered saline
AVI	audio video interleave	PC	phase contrast
BSA	bovine serum albumin	PDZ	PSD95 DlgA zo-1
CA1	<i>Cornu Ammonis</i> area 1	PFA	paraformaldehyde
CCD	charge-couple device	PKCα	protein kinase C alpha
cDNA	complementary deoxyribonucleic acid	PNS	peripheral nervous system
ChAT	choline acetyltransferase	PSD	postsynaptic density
CNS	central nervous system	QD	quantum dot
COS cells	kidney cell line from the African green monkey	QDOT	quantum dot
D	diffusion coefficient	RNA	ribonucleic acid
DA	dopamine; 4-(2-aminoethyl)benzene-1,2-diol	RT	room temperature
DIC	differential interference contrast	SEM	standard error of the mean
DIV	days <i>in vitro</i>	SA	self-administration
DMSO	dimethyl sulfoxide	SFK	src-family kinases
DNA	deoxyribonucleic acid	shRNA	short hairpin ribonucleic acid
DNase	deoxyribonuclease	SPT	single particle tracking
E18	embryonic day 18	SV	Sindbis virus
EDTA	ethylene diamine tetraacetic acid	TM	transmembrane
EGFP	enhanced green fluorescent protein	TTX	tetrodotoxin
EGTA	ethylene glycol tetraacetic acid	VDCC	voltage-dependent Ca <sup>2+</sup> channels
EM	electron microscopy	VDSC	voltage-dependent Na <sup>+</sup> channels
ER	endoplasmic reticulum	VGAT	vesicular γ-aminobutyric acid transporter
EYFP	enhanced yellow fluorescent protein	vGlut	vesicular glutamate transporter
FCS	fetal calf serum	VIAAT	vesicular inhibitory amino acid transporter
FRAP	fluorescence recovery after photobleaching	VTA	ventral tegmental area
GABA	γ-aminobutyric acid	WT	wild type
GABA <sub>A</sub> R	γ-aminobutyric acid type A receptor	YTH	yeast two-hybrid
GAD	glutamate decarboxylase		
GFP	green fluorescent protein		
Glu	glutamate		
GluR	glutamate receptor		
Gly	glycine		
GlyR	glycine receptor		
GST	glutathione S-transferase		
HA	anti-hemagglutinin		
HEK cells	Human Embryonic Kidney 293 cells		
His-tag	hexa histidine-tag		
HRP	horse radish peroxidase		
IgG	immunoglobulin G		
IP	immunoprecipitation		
IPSC	inhibitory postsynaptic current		
IPTG	Isopropyl β-D-1-thiogalactopyranoside		
IR	immunoreactivity		
IU	international unit		

---

## Summary

In the brain,  $\alpha 7$  neuronal nicotinic acetylcholine receptors ( $\alpha 7$  nAChRs) have a special role amongst nAChRs.  $\alpha 7$  nAChRs are forming homopentamers, they display a high permeability for  $\text{Ca}^{2+}$ , and they are the most prevalent nAChRs in the brain.  $\alpha 7$  nAChRs are found at the highest concentration in the hippocampus where they are located mostly on GABAergic interneurons and play an important role in learning and memory. Moreover  $\alpha 7$  nAChRs have been involved in diseases such as Alzheimer's disease (AD) and schizophrenia, and are attracting considerable scientific interest to elucidate their contribution to disease mechanisms. While central cholinergic circuits have been investigated very extensively, the cell- and molecular biological properties of  $\alpha 7$  nAChRs have not been studied in depth. The exact subcellular localization of  $\alpha 7$  nAChRs is still debated, in particular in relation to synaptic sites, and only two proteins interacting with  $\alpha 7$  nAChR, namely RIC-3 and Src-family kinases, have been identified to date. However, none of them is involved in synaptic clustering of  $\alpha 7$  nAChR. In chapter 2 we describe the discovery of PICK1 as a novel  $\alpha 7$  nAChR interacting protein. Thereby the PDZ domain of PICK1 binds to the large cytoplasmatic loop of  $\alpha 7$  nAChR. We present evidence that PICK1 regulates clustering of  $\alpha 7$  nAChRs in rat hippocampal interneurons.

The more detailed investigation of  $\alpha 7$  nAChR clustering and surface dynamics demanded the ability to express exogenous gene constructs in cultured neurons. In chapter 3 we report an optimized transfection protocol for rat hippocampal neurons. Use of the magnetofection technique allowed the parallel transfection of several constructs and their expression in neurons for up to 3 weeks *in vitro*.

To understand the clustering and localization of  $\alpha 7$  nAChRs it is necessary to investigate the surface dynamics of single receptors. In chapter 4 we report a detailed analysis of  $\alpha 7$  nAChR cell surface mobility, using  $\alpha$ -BT and QDots labeled single receptor trafficking.  $\alpha 7$  nAChRs were found to be very mobile within the membrane. Clusters were found to be mobility traps, suggesting  $\alpha 7$  nAChRs interact with underlying scaffolding proteins at these sites. Mobility

---

traps were found extrasynaptically and perisynaptically in close vicinity to GABAergic and glutamatergic postsynaptic densities. While extrasynaptic  $\alpha 7$  nAChRs might activate  $\text{Ca}^{2+}$ -dependent signaling pathways, the perisynaptic  $\alpha 7$  nAChRs are probably playing a modulatory role in GABAergic and glutamatergic synaptic activity.  $\alpha 7$  nAChR mobility was not only dependent on localization but also on chronic synaptic activity changes and activation of the receptor itself.

Taken together, in this thesis work we identify  $\alpha 7$  nAChR as a highly regulated receptor. The sites of  $\alpha 7$  nAChR-dependent  $\text{Ca}^{2+}$  influx are tightly controlled by the cell.  $\alpha 7$  nAChRs are clustered at distinct sites, reflecting functional heterogeneity. We identify for the first time a direct protein-protein interaction mechanism involved in the regulation of  $\alpha 7$  nAChR clustering and possibly surface expression. We uncover  $\alpha 7$  nAChR clusters as sites where mobility is constrained, but single receptors are able to diffuse in and out, confirming receptor clusters as steady-state receptor aggregations. We find  $\alpha 7$  nAChR distributed all over the cell surface with clusters formed at extra- and perisynaptic sites. We speculate that  $\alpha 7$  nAChRs have a variety of different functions dependent on their localization. The distinct mechanisms of the particular  $\alpha 7$  nAChR subpopulations remain unclear, and are left to be addressed in future work.

---

## Zusammenfassung

Alpha7 neuronale nikotinische Azetylcholinrezeptoren ( $\alpha 7$  nACh Rezeptoren) unterscheiden sich im Gehirn von anderen nikotinischen Azetylcholinrezeptoren durch mehrere Eigenschaften. Zum Einen sind  $\alpha 7$  nACh Rezeptoren die häufigsten Azetylcholinrezeptoren im Gehirn. Zum Anderen sind die homopentameren Rezeptoren äusserst permeabel für Kalziumionen. Die höchste Dichte von  $\alpha 7$  nACh Rezeptoren finden wir im Hippocampus auf GABAergen Interneuronen wo sie eine wichtige Rolle in Lern- und Gedächtnisprozessen spielen. Weiter wurde der  $\alpha 7$  nACh Rezeptor als Akteur in verschiedenen Krankheiten, wie Alzheimer-Krankheit und Schizophrenie, identifiziert. Dies hat zu vermehrten Anstrengungen für die Aufklärung der Funktionsweise des Rezeptors geführt. Während zentrale cholinerge Vernetzungen ausführlich untersucht wurden, ist die Bedeutung von  $\alpha 7$  nACh Rezeptoren auf Zell- und Molekularebene weitestgehend unklar. Die genaue subzelluläre Lokalisierung von  $\alpha 7$  nACh Rezeptoren, speziell in Hinsicht auf die synaptische Umgebung, ist bis heute umstritten. Zudem wurden bis heute nur zwei Interaktionsproteine von  $\alpha 7$  nACh Rezeptoren identifiziert. Jedoch scheint keines der beiden Interaktionsproteine, RIC-3 und Kinasen aus der Src-Familie, einen Einfluss auf die synaptische Aggregation von  $\alpha 7$  nACh Rezeptoren zu haben. In Kapitel 2 beschreiben wir die Entdeckung von PICK1 als ein weiteres interagierendes Protein. Wir zeigen, dass die Interaktion über die PDZ Domäne von PICK1 und dem grossen zytoplasmatischen Teil des  $\alpha 7$  nACh Rezeptors abläuft. Zudem zeigen wir, dass PICK1 die Aggregation des  $\alpha 7$  nACh Rezeptors in hippocampalen Interneuronen aus der Ratte reguliert.

Eine ausführliche Untersuchung der Aggregation und der Oberflächendynamik des  $\alpha 7$  nACh Rezeptors verlangt nach der Möglichkeit fremde Gensequenzen in kultivierten Neuronen zu exprimieren. In Kapitel 3 beschreiben wir ein solches optimiertes Transfektionsprotokoll für hippocampale Neuronen aus der Ratte. Die Verwendung der Magnetofektionstechnik erlaubte uns die parallele Transfektion und



---

Expression für bis zu 3 Wochen von mehreren Konstrukten in *in vitro* kultivierten Neuronen.

Um die Aggregation und die Lokalisation des  $\alpha 7$  nACh Rezeptors zu verstehen, ist es notwendig die Oberflächendynamik eines einzelnen Rezeptors zu untersuchen. Im Kapitel 4 berichten wir in einer ausführlichen Analyse von der Zelloberflächenmobilität des  $\alpha 7$  nACh Rezeptors. Dabei wurden einzelne Rezeptoren mit  $\alpha$ -BT und QDots markiert. Wir identifizierten den  $\alpha 7$  nACh Rezeptor als sehr mobilen Rezeptor innerhalb der Membran. Aggregationen von  $\alpha 7$  nACh Rezeptoren kommen durch lokale Mobilitätsverringerng des Rezeptors zustande, was eine Interaktion zu zytosolischen Gerüstproteinen vermuten lässt. Solche Effekte konnten wir in extrasynaptischen wie auch perisynaptischen Stellen, nahe zu GABAergen und glutamatergen postsynaptischen Dichten, finden. Während die extrasynaptischen  $\alpha 7$  nACh Rezeptoren wahrscheinlich kalziumabhängige Signalkaskaden aktivieren, haben die perisynaptischen  $\alpha 7$  nACh Rezeptoren möglicherweise eine modulatorische Funktion. Zusätzlich zur Lokalisation hat auch die chronische synaptische Aktivität und die Aktivierung des Rezeptors selbst einen Effekt auf die Oberflächenmobilität.

Zusammengefasst, in dieser Arbeit identifizieren wir den  $\alpha 7$  nACh Rezeptor als stark regulierten Rezeptor. Die Lokalisation des Rezeptors und somit der rezeptorabhängige Kalziumeinstrom wird durch die Zelle streng kontrolliert. Die unterschiedlichen Lokalisationen des Rezeptors lassen eine funktionale Heterogenität des  $\alpha 7$  nACh Rezeptors vermuten. Wir beschreiben erstmalig eine Proteininteraktion mit dem  $\alpha 7$  nACh Rezeptor welche die Rezeptoraggregation und dessen Oberflächenexpression reguliert. An Stellen wo die Rezeptormobilität stark eingeschränkt ist, beobachten wir nACh Rezeptoraggregationen. Einzelne Rezeptoren diffundieren jedoch in und aus der Aggregationstelle, so dass die Aggregation statt als fixes Gebilde besser als stetiger Zustand beschrieben werden sollte. Wir finden  $\alpha 7$  nACh Rezeptoren diffus und aggregiert in extrasynaptischen und perisynaptischen Stellen. Wir vermuten, dass  $\alpha 7$  nACh Rezeptoren je nach Lokalisation unterschiedliche Funktionen wahrnehmen. Die Aufklärung dieser Funktionen wird Teil zukünftiger Forschung sein.

## 1. Introduction

Neurotransmitter receptors are fundamental components of neuron-neuron communication. The localization and density of neurotransmitter receptors on neurons in the central and the peripheral central nervous system (CNS, PNS) are tightly regulated. Neurotransmitter receptors are aggregated in clusters at sites of their function. Any dysfunction in this regulation leads to neurological adverse consequences as occurring in several diseases. In this thesis we focused on the  $\alpha 7$  neuronal nicotinic acetylcholine receptor ( $\alpha 7$  nAChR). We investigated the clustering and localization of  $\alpha 7$  nAChR in neuronal hippocampal cultures. We identified binding partners of  $\alpha 7$  nAChR and their effect on  $\alpha 7$  nAChR clustering. By using single particle tracking methods we further shed light on the cell surface dynamics of single  $\alpha 7$  nAChR in respect to the localization of  $\alpha 7$  nAChR cluster as well as synapses.

### 1.1. Discovery of the acetylcholine receptor

#### 1.1.a. Discovery of chemical synaptic transmission


The  $\alpha 7$  nAChR is a member of the family of the nicotinic acetylcholine receptors. nAChRs played an important role in the historical discovery of chemical neurotransmission. After decades of wild speculations and theories about the working mechanism of nerve and muscle in the Renaissance, the work of Galvani and Matteucci elucidated for the first time that the nerve-muscle signal transduction is based on electrical propagation (Matteucci, 1842). Based on this finding Claude Bernard could show in 1850 by using curare that the electrical impulses are transferred from the nerve to the muscle (Bernard, 1850). Just a few years later, in 1866, Vulpain suggested that the nerve and the muscle are not one unit. He proposed the existence of a neuromuscular junction (NMJ) between the nerve and the muscle (Vulpain, 1866). How the signal from the nerve overcomes this gap remained unclear until John Langley was investigating the effect of nicotine and curare 1905. Based on his findings he suggested that a transmitter-substance is released at the nerve terminal, propagating the signal to the muscle (Langley, 1905). Still the natural ligand was unclear. By 1914 Henry Dale isolated a compound

secreted by fungus ergot (*Claviceps*), which he identified as acetylcholine (ACh) (Dale, 1914b). He thoroughly investigated the actions of ACh and found that it produced two major responses: a 'muscarine' and a 'nicotine' action. Muscarine was isolated in 1869 by Schmiedeberg and Koppe from *Amanita muscaria*. They showed that muscarine arrests the beat of the frog heart (Schmiedeberg and Koppe, 1869). Dale described that the muscarine actions of ACh were antagonized by atropine, whereas the nicotine actions were antagonized by curare. Influenced by the work of his colleague Otto Loewi, he could show by extensive studies that ACh is a neurotransmitter on the smooth muscle cell but also at synapses at the autonomic ganglia. For the 'discovery of chemical synaptic transmission' he and Otto Loewi were awarded with the Nobel Prize in Physiology and Medicine in 1936.

### **1.1.b. Isolation and characterization of the nAChR**

After the identification of ACh as a neurotransmitter tremendous work has been done to identify competitors of ACh. The presence of an acetylcholine receptor protein was speculated (Nachmansohn, 1955). But only with the discovery of  $\alpha$ -bungarotoxin, a cholinergic antagonist isolated from the snake venom from *Bungarus multicinctus* (Chang and Lee, 1963), Changeux et al. finally managed in 1970 to confirm the existence of an ACh receptor in preparations of the electric organ of *Electrophorus electricus* (Changeux et al., 1970). According to the binding properties of nicotine, the receptor was termed nicotinic acetylcholine receptor. Further progress in the isolation of nAChRs of *Electrophorus electricus* (**Fig. 1-1**) (Olsen et al., 1972) allowed to propose that the nAChR is a pentameric protein (**Fig. 1-2**) (Hucho and Changeux, 1973). With the development of new techniques such as reverse genetics and molecular cloning the pentameric *Torpedo californica* nAChR was found to consist of related but genetically distinct subunits (**Fig. 1-1**). The subunits were classified according to similarities in protein sequences in  $\alpha$ ,  $\beta$ ,  $\delta$ ,  $\varepsilon$  and  $\gamma$  subfamilies (Noda et al., 1982; Claudio et al., 1983; Noda et al., 1983a; Noda et al., 1983b; Takai et al., 1985). In contrast to non- $\alpha$  subunits the  $\alpha$  subunit comprises a Cys-Cys pair at the extracellular side near to the transmembrane region 1 (TM1) which is crucial for ligand binding (Karlin et


al., 1986). Screening of brain cDNA libraries led to the discovery of several closely related sequences which encode 16 structurally homologous subunits. Further the subunits were subdivided into muscle or neuronal subunits. In muscle only two subunit compositions of nAChRs were discovered, that are  $2\alpha 1, \beta 1, \gamma, \delta$ , and  $2\alpha 1, \beta 1, \epsilon, \delta$ . Whereas in neuronal tissue several  $\alpha$  subunits ( $\alpha 2 - \alpha 7$  and  $\alpha 9 - \alpha 10$ ;  $\alpha 8$  has been discovered in avian libraries but not in mammals) and  $\beta$  subunits ( $\beta 2 - \beta 4$ ) were identified (Le Novere et al., 2002). In contrast to the muscle AChRs the neuronal AChR subunits form more versatile pentamers.  $\alpha 7$ - $\alpha 10$  exist each as homopentamers or as  $\alpha 7/\alpha 8$  and  $\alpha 9/\alpha 10$  heteropentamers.  $\alpha 2$ - $\alpha 6$  and  $\beta 2$ - $\beta 4$  are included in a range of complex heteromers (Le Novere et al., 2002).



**Fig. 1-1 Discovery of nAChR:** The nAChR was first isolated from the electric organ of *Electrophorus electricus*. Another source of nAChR is the electric tissue of *Torpedo californica*.

One has to note that even though most of the neuronal nAChR subunits are expressed in neurons, they are also found in some non-neuronal cells throughout the body (Kawashima and Fujii, 2003; Gahring and Rogers, 2005). Further it is important to mention at this point that apart from nicotinic AChRs another group of ACh receptors exist, muscarinic acetylcholine receptors (mAChR) named after their binding affinity of muscarine. In contrast to nAChRs, which are ligand gated ion channels, the mAChRs are G protein coupled receptors which have slower and longer lasting effects. mAChRs activate downstream signaling via activation of phosphoinositide-specific phospholipase C $\beta$  and Ca<sup>2+</sup> release from internal Ca<sup>2+</sup> stores (Eglen, 2006). Moreover, mAChRs are also involved in inhibition of elevated adenylate cyclase activity.

The studies with *Torpedo californica* nAChRs revealed the structural characteristics which are relevant to all nAChR subunits identified to date. Generally, all nAChRs assemble in a pentameric arrangement forming an internal membrane spanning pore which is permeable to a different degree for  $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Ca}^{2+}$ . Each subunit consists of 4 TM regions with the 3<sup>rd</sup> and 4<sup>th</sup> TM region being linked by a large cytoplasmic loop. The N- and the C-terminals are extracellular (**Fig. 1-2**). Though the high structural similarities of the different subunits, nAChRs formed by the set of  $\alpha$ ,  $\beta$ ,  $\delta$ ,  $\gamma$ ,  $\epsilon$  subunits feature great differences in ligand affinity (or anta-, agonist affinity), ion permeability, ion selectivity, and desensitization rate (Le Novere et al., 2002). Tissue specific nAChR subunit expression ensures adapted function of cholinergic receptors dependent on their localization.




**Fig. 1-2 Architecture of the nAChR:** Ribbon diagrams of the whole *Torpedo californica* acetylcholine receptor at 4Å resolution. Viewed (**A**) from the synaptic cleft and (**B**) parallel with the membrane plane. In (**A**) the transmembrane pore formed by the 5 subunits is clearly visible. For clarity, only the ligand-binding domain is highlighted in (**A**) and only the front two subunits are highlighted in (**B**). The grey bars in (**B**) indicate the membrane. Schematic drawing of a single subunit (**C**). The subunit spans the membrane 4 times. The 3<sup>rd</sup> and 4<sup>th</sup> TM element are connected by a large cytoplasmic loop. The C- and N-terminal are extracellular. The ligand binding site is located at the N-terminal of the  $\alpha$ -subunits. (Unwin, 2005)

## 1.2. Principles of neurotransmission

### 1.2.a. From action potential to muscle contraction

The discovery of the neuromuscular junction (NMJ) as the first observed synapse, the relative large size of the synaptic apparatus and the well established purification methods were reasons for the NMJ to become a

model system in synapse research. In contrast to synapses in the central nervous system (CNS) the NMJ connects two very different types of cell types, namely neuronal cells and muscle cells. Nevertheless the NMJ is a perfect system to illustrate the principles of neurotransmission. As in all types of synapses the NMJ harbors the key players of a synaptic contact. Basically, a synapse propagates an electric signal, an action potential (AP), from the presynapse to the postsynapse of a connected cell. Therefore the electrical signal is transformed at the cell-cell junction into a chemical signal by the release of neurotransmitters from the active zone into the narrow synaptic cleft. The neurotransmitters are detected in the neighboring cell by neurotransmitter receptors in the postsynaptic density (PSD). The PSD is a protein network harboring all components of the postsynaptic apparatus. Activation of the neurotransmitter receptor results in the opening of the internal pore and the influx of ions. The inward directed current results in the generation of an AP and propagation of the electric signal. At the NMJ the arriving AP opens presynaptic voltage-dependent calcium channels and  $\text{Ca}^{2+}$  flows into the motor neuron's presynaptic cytosol. The presynaptic  $\text{Ca}^{2+}$  influx triggers the fusion of neurotransmitter containing vesicles with the membrane eliciting the release of ACh into the synaptic cleft. ACh diffuses to the motor endplate where it specifically binds to nAChRs. nAChRs, which are ligand-gated ion channels, open and allow  $\text{Na}^+$  and  $\text{Ca}^{2+}$  influx according to their gradient from the extracellular lumen into the cytosol of the muscle cell. The ion influx leads to a depolarization of the muscle cell. The depolarization further opens voltage gated sodium channels. The influx of  $\text{Na}^+$  ions generates an action potential which spreads within the muscle cell. Consequently the sarcoplasmic reticulum releases  $\text{Ca}^{2+}$ , eliciting the contraction of the muscle (**Fig. 1-3**).




**Fig. 1-3 Structure of the neuromuscular junction:** Typical for the NMJ is the big terminal and the folded postsynaptic membrane. **(1)** An AP is propagated from the axon to the presynaptic terminal. **(2)** Voltage-dependent  $\text{Ca}^{2+}$  channels (VDCC) open.  $\text{Ca}^{2+}$  enters the presynaptic cytosol. **(3)** Primed synaptic vesicles, filled with ACh, fuse with the cell membrane. ACh is released into the synaptic cleft. **(4)** AChRs open upon ACh binding and allow  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  influx and membrane depolarization. **(5)** Voltage-dependent  $\text{Na}^{+}$  channels open. **(6)**  $\text{Na}^{+}$  influx induces  $\text{Ca}^{2+}$  release from the sarcoplasmic reticulum which activates muscle contraction. Adapted in part from (Kandel et al., 2000)

### 1.2.b. NMJ vs. neuronal synapse

Even though the NMJ is a very useful system to investigate the architecture of a synapse and to uncover the molecular principles of neurotransmission, the system is limited. The NMJ differs in many aspects from a synaptic contact between neurons in the central nervous system (CNS). In comparison to the NMJ a neuronal synapse is around 1000 times smaller. The postsynaptic density (PSD) of the NMJ is folded several times to increase the surface allowing a higher receptor number in the synaptic cleft whereas the neuronal PSD has a small and unfolded surface and harbors just a fraction of the NMJ receptor number. In the folded synaptic cleft of the NMJ resides the basement membrane, or basal lamina (**Fig. 1-3**). A comparable structure in the neuronal synaptic cleft is absent (**Fig. 1-4**). The enzyme acetylcholine esterase (AChE), which degrades ACh to choline and acetate is aggregated in the basal lamina. While the NMJ PSD comprises only two types of nAChRs, the neuronal PSD can be excitatory or inhibitory, harboring glutamate receptors (GluR) and GABA receptors (GABAR), respectively. Accordingly, the neuronal synapses feature various types of presynapses, whereas the uniform NMJ presynapse stores only ACh-containing vesicles. In regard to the varieties between the two classes of synapses it is evident that the underlying PSD machinery is totally different. The differences between the NMJ and neuronal synapses reflect their functional specialization. The synaptic contact at the muscle end plate is optimized to propagate and amplify a single AP in the muscle with maximal reliability. In the CNS the focus lies less on the transmission of a single AP but rather on integration of multiple signals within the neuronal network and on the complexity of the network itself. In contrast to the simple architecture of motor neurons, the complex organization of neuronal networks demands regulatory mechanisms orchestrating excitatory actions.





**Fig. 1-4 Structure of a neuronal synapse:** In contrast to the NMJ the postsynaptic membrane is not folded. Though, the mechanism of signal transduction is similar. **(1)** An AP arrives at the presynaptic terminal. **(2)** Voltage-dependent  $\text{Ca}^{2+}$  channels (VDCC) open and  $\text{Ca}^{2+}$  enters the cell. **(3)**  $\text{Ca}^{2+}$  triggers fusion of the neurotransmitter-filled synaptic vesicles with the plasma membrane. **(4)** Neurotransmitters enter the synaptic cleft. Upon binding of neurotransmitters to the neurotransmitter receptors, ions enter the postsynaptic cytosol. **(5)** Neurotransmitters are released from the receptor and diffuse away from the synaptic cleft where they get degraded or recycled by neurotransmitter re-uptake pumps.

### 1.2.c. An evolutionary point of view

During evolution from primitive animals with a simple neuronal network, such as jellyfish or worms, to species with a more complex and larger neuronal network and brain, the necessity for regulation of synaptic activity became more and more important. In early ancestor animals synaptic contacts were mainly excitatory whereas more complex species developed neuronal inhibitory mechanisms and synchronized neuronal networks to protect the brain from overexcitation. It has been shown that the proportion of cortical

inhibitory neurons correlates with the size of the brain (Glendenning, 1998). In the human brain inhibitory interneurons comprise 15-25% of the cortical neurons (Markram et al., 2004). Computational simulations and *in vivo* experiments brought evidence that the interplay of excitation and inhibition is extremely important to ensure the proper function of the brain (Dichter and Ayala, 1987; Vogels et al., 2005). A loss of inhibition leads to harmful epileptic activity.

In the course of evolution an ancestor ligand-gated cation permeable neurotransmitter, emerged through an early lateral transfer from a prokaryotic source 540 million years ago. By gene duplications and mutations the ancestor gave rise to a set of paralogs of structurally similar but functionally different neurotransmitter receptors (Tasneem et al., 2005). The paralogue receptors are oligomers forming a channel pore through the membrane allowing the selective influx of ions into the cytosol. Though the receptors exhibit different ion selectivity and are sensitive to different ligands their main structure is conserved supporting the hypothesis of a common origin. GABA<sub>A</sub>R, nAChR, 5-HT<sub>3</sub> and GlyR are all arranged as a pentamer with subunits harboring 4 TM regions. NMDAR, AMPAR and kainate receptor differ slightly from the pentameric receptors. Their subunits span the membrane 4 times but are arranged as a tetramer. In contrast to the structural similarities the functional differences are more pronounced among the set of ligand-gated ion channels. While NMDAR, AMPAR and kainate receptors are permeable for cations inducing excitation in a neuron, GABA<sub>A</sub>R and GlyR are inhibitory by allowing Cl<sup>-</sup> ions to enter the cell and increasing the AP threshold level by hyperpolarization.

#### **1.2.d. Architecture of a neuronal synapse**

A theoretical neuronal network is built of a high number of neuronal units which are connected.

In the CNS a single neuron gets input from several thousands of other neurons. They release their neurotransmitters on synaptic receptors localized mainly on dendrites, causing excitation or inhibition. Excitation in the receiving neuron further results in the axonal release of cell type specific neurotransmitters onto hundreds of connected neurons. It is evident that a


complex neuronal network must be structurally organized and functionally tightly regulated. Therefore the arrangement and strength of synaptic contacts are crucial. In principal neurons of the neocortex the excitatory and inhibitory input on the dendrite is spatially separated (**Fig. 1-5**). Excitatory synaptic input from an axon is received on a dendritic membrane protrusion containing the postsynaptic machinery. These protrusions are called spines (**Fig. 1-6**). The dendrites of a single neuron possess from thousands up to a few hundred thousand spines. Spines incorporate the complete excitatory postsynaptic machinery.

The components of this machinery are centralized in the PSD. The glutamatergic PSD comprises dozens of different proteins with cytoskeletal and regulatory functions. PSD-95, Homer, Grip, and PICK1 are just a few of the many proteins playing a role in the structural matrix holding ion channels, kinases, and phosphatases in place, on the tip of the spine apposed to the active zone of the presynapse (**Fig. 1-5**). Spines maintain existing and establish new synaptic contacts with innervating axons.

Inhibitory postsynapses are fewer in number. The inhibitory PSD is in immediate proximity to the excitatory PSDs on the dendritic shaft of the same dendrite. There it receives axonal inputs from GABAergic or glycinergic axon terminals (**Fig. 1-5**). The close proximity of excitatory and inhibitory synapses requires a tight control of the localization of all synaptic proteins. The proper functionality of a synapse demands further that pre- and postsynapse are well aligned with a defined distance. Inconsistencies in the architecture of the synaptic cleft would affect the speed and probability of the AP induction. Additionally, the amount of released neurotransmitters and the number of opposing neurotransmitter receptors need to be balanced for synaptic transmission. To ensure that an AP and the following neurotransmitter release into the synaptic cleft overcomes the postsynaptic depolarization threshold level and provokes the generation of an AP in the postsynaptic cell, the number of responsive neurotransmitter receptors in the synaptic cleft has to reach a certain number. In addition to the receptor number the receptor localization has to be strictly regulated. Released neurotransmitters diffuse a few hundred nanometers in the synaptic cleft before they get enzymatically degraded or recycled. Therefore neurotransmitter receptors need to be

localized in immediate vicinity to the neurotransmitter release site. Receptor number and localization are two parameters which are tightly regulated by the postsynaptic cell. An imbalance in this regulation is known to cause diseases such as depression, schizophrenia, and addiction (Nader et al., 2006; Witkin et al., 2007; Kessler et al., 2009). The overall receptor number is adjusted by the rate of protein synthesis, rate of internalization and integration from and to the membrane, respectively. However, the receptor number in the PSD is dependent on specific anchoring mechanisms keeping the receptor at its place of function. The regulation of receptor number in the PSD goes in hand with the control of receptor localization. It has been shown that receptors are stabilized and anchored through different mechanisms (Allison et al., 1998). Receptors can be anchored directly or via an adaptor protein to cytoskeletal protein scaffolds. The actin cytoskeleton or the microtubule network provide the stability for receptor anchoring (**Fig. 1-5**). Another anchoring mechanism excludes cytoskeletal protein networks but consists of membrane patches with an interlaced protein network providing a scaffold for receptor stabilization.

Altogether, synaptic transmission at the synapse is dependent on many factors. Besides the complex presynaptic machinery guaranteeing tuned neurotransmitter release, the PSD is responsible for adjusted receptor number and localization which involves membranous or cytoskeletal scaffolding proteins interacting directly or indirectly with neurotransmitter receptors.



**Fig. 1-5 Excitatory and inhibitory innervation:** (left)

Excitatory input is received from Glutamate (Glu) releasing terminals. Glutamate receptors (GluR) as AMPA receptors (AMPA) and NMDA receptors (NMDAR) are located in spines. Actin ensures the shape of the spine while the postsynaptic density (PSD) protein network stabilizes the neurotransmitter receptors. Activation of AMPARs and NMDARs by Glu leads to  $\text{Ca}^{2+}$  and  $\text{Na}^{+}$  influx into the postsynaptic cytosol. (right) Inhibitory innervation takes place at the shaft of the dendrite. GABA or Glycine (Gly) is released into the synaptic cleft from the presynaptic terminal upon arrival of an AP.  $\text{GABA}_A$  receptors ( $\text{GABA}_A\text{R}$ ) and Glycinergic receptors (GlyR) are stabilized by Gephyrin and further unknown proteins. Following activation of  $\text{GABA}_A\text{Rs}$  or GlyRs,  $\text{Cl}^{-}$  enters and hyperpolarizes the cell.


### 1.3. From static to dynamic view

#### 1.3.a. The tools for microscopy

In the past, all insights into the architecture of the NMJ and synaptic contacts in the brain were derived from microscopy studies. Thereby the tissue was stained with different dyes enhancing the contrast of tissue structures. Years later, antibody labeling further allowed elucidating the distribution of distinct proteins.

In the 19<sup>th</sup> and early 20<sup>th</sup> century stained tissue sections were mainly investigated with optical microscopes. The resolution of these analyses was limited due to the physical restriction of the wavelength of visible light. With the invention of the electron microscope in 1931 by Ernst Ruska, who was awarded with the Nobel Prize in Physics in 1986 for his discovery, biological samples could be investigated on an ultrastructural level. This fascinating tool

allowed the observation of single proteins and even lipid bilayers of the plasma membrane and vesicles (**Fig. 1-6**).



**Fig. 1-6 Neuronal synapse viewed at with electron micrograph (EM):** EM allows observations on an ultrastructural level. Darker areas are lipid rich structures whereas lighter areas are devoid of lipids. The image illustrates an excitatory synapse in the hippocampus. The large synaptic terminal is filled with synaptic vesicles (**blue**) and harbors one mitochondrion (**red**). Two postsynaptic spines are opposed (**green**). The PSD is clearly visible as dark band facing the synaptic cleft. Scale bar: 500 nm (Source: Caroline Petitjean)

Even though these methods reveal the structural architecture of cells, organelles, and also synapses, in an unprecedented detail-rich manner, the data just allow a static interpretation. This limitation motivated scientists to develop methods to collect in addition to spatial data also temporal data. The temporal visualization of a biological system provides important information about the dynamic processes taking place, providing a closer look at the real situation. The first live microscopy experiments were performed on large size specimens such as plant or animal cells without visualizing certain structures in the living cell. Deeper insights into cell processes such as protein localization, turnover and diffusion, or the dynamics of intracellular compartments such as endosomes became feasible with the discovery of the green fluorescent protein (GFP) and its use as a marker for protein expression (Chalfie et al., 1994). GFP is a fluorescent protein, which when exposed to blue light emits in the green spectra. It was isolated from jellyfish

(*Aequorea victoria*) in 1962 by Osamu Shimomura (Shimomura et al., 1962). In 2008 he and his colleagues Martin Chalfie and Roger Y. Tsien were awarded with the Nobel Prize in Chemistry for 'the discovery and development of the GFP'. GFP allows the detection of localization and movement of a protein of interest in living cells and even animals. Therefore, GFP is linked to the protein of interest by molecular cloning and expressed in cell culture or tissue. In the past years GFP was developed further and optimized, providing today a palette of different coloured fluorescent proteins and even a pH sensitive GFP. The use of GFP and its derivatives as markers in living cells established a new era of live microscopy research.

### **1.3.b. The brain – a dynamic structure**

Live microscopy experiments with living neurons changed the understanding of synaptic contacts. With the acquisition of microscopy data in living neuronal tissue over a longer time period it became obvious that synaptic contacts are not rigid and inflexible structures, but rather that the neuronal network is very dynamic and underlies a constant change, even in the aged brain. Using two-photon microscopy and confocal microscopy dendritic spines were identified as very dynamic structures which change their shape within minutes and build up or brake synaptic contacts in the range of hours (Dunaevsky et al., 1999). These "plastic" changes in spine morphology and synaptic contact formation are mostly activity-dependent, displaying an adaptation mechanism reacting to neuronal activity or external input. Synaptic plasticity is believed to underlie motivation, learning, and memory (Hebb, 1949). On a closer look, synaptic plasticity is the strengthening or weakening of single synaptic contacts. This is regulated pre- or postsynaptically. Presynaptically, the modulation of the synaptic strength is achieved by regulation of the amount of neurotransmitter release per arriving action potential (Debanne et al., 2003). Postsynaptically, synaptic strength is controlled by the number and properties of neurotransmitter receptors (Debanne et al., 2003). The development of new tools to investigate single receptors within the membrane gave deeper insights into the membrane dynamics and the diffusion behaviour of receptors. Thereby single receptors are labelled with small fluorescent probes, composed of a semiconductor material. These probes, known as

quantum dots (QDOTs), have a remarkable photo stability and high quantum yield (Chan and Nie, 1998). These two features permit the detection and recording of single receptors over a long time period using a light microscope. From the recorded trajectories, parameters including diffusion coefficient, confinement, and mean square displacement can be determined (Bannai et al., 2006). Earlier studies labelled single receptors with silicon beads which are much larger in size compared to QDOTs (Meier et al., 2001). In contrast to silicon beads, QDOT labelling of receptors does not alter the diffusion behaviour within the membrane. Slight changes of receptor diffusion were only observed within the synaptic cleft. Even though QDOT coupled receptors are able to enter the synaptic cleft, the freedom of mobility is reduced due to the bigger size of the QDOT-antibody-receptor complex (Groc et al., 2007; Triller and Choquet, 2008).

Single particle tracking experiments of different labelled neurotransmitter receptors like GlyR, AMPAR, and NMDAR revealed a highly dynamic diffusion within the plasma membrane (Dahan et al., 2003; Groc et al., 2004). Neurotransmitter receptors are accumulated and clustered at the postsynaptic density. In addition single receptors diffuse freely within the plasma membrane. These findings suggest that receptor clusters are sites where receptors are slowed down by interacting with membrane proteins or cytosolic scaffolding proteins. In view of the fact that single receptors are just trapped temporary within a cluster points to a steady-state model in which receptors are constantly exchanged with the surrounding. Apart from lateral membrane diffusion, receptor number and localization is also dependent on dynamic intracellular processes. After a receptor is synthesised at the endoplasmatic reticulum (ER) and processed at the Golgi complex it is transported to endosomes from where it is further directed to the membrane. There, neurotransmitter receptors, diffuse freely within the membrane and eventually get confined in synaptic sites where they are activated by presynaptically released neurotransmitters. In parallel to receptor integration, receptors are constantly internalized from the plasma membrane and either recycled via endosomes or degraded within lysosomes.

With the development of new tools and microscopes it became evident in the last decade that neurotransmitter receptors are not statically integrated in the




membrane and synaptic contacts are not rigid constructs. Instead postsynaptic proteins such as scaffolding proteins and neurotransmitter receptors are dynamically regulated and are constantly in motion. These findings resulted in the view of the brain as a highly adaptive and a permanently altering neuronal network. Additionally, the understanding and the treatment of several diseases caused by an imbalance in the synaptic composition are now better understood from a dynamic-model point of view.

## **1.4. Nicotinic acetylcholine receptors in the brain**

### **1.4.a. The cholinergic system**

nAChRs are present on autonomic neurons, namely preganglionic neurons of the parasympathicus, on adrenal chromaffin cells in the PNS, on motoneurons in the NMJ and on many neurons in the CNS. The nAChRs in the CNS are classified as neuronal nAChRs. The diversity of nAChR localization implies the different roles in cholinergic transmission. Cholinergic systems are involved in arousal, sleep-wakefulness, locomotor behaviour, learning and memory (Woolf, 1991). The neuronal nAChRs became of interest in the recent past. Before that time, the focus was on the main excitatory and main inhibitory circuits in the brain. The glutamate receptors AMPAR and NMDAR are the most prominent receptors involved in excitation while GABA is the main inhibitory neurotransmitter in the mammalian brain. In contrast, cholinergic circuits comprise just a minority of all circuits (Woolf, 1991). The finding of the involvement of cholinergic processes in diseases such as schizophrenia in the last years resulted in increased effort in the uncovering of the cholinergic system. This was supported by the development of sensitive immunohistochemical methods, using choline acetyltransferase (ChAT) as a marker for cholinergic cells. In addition to motoneurons, there are three major cholinergic subsystems that innervate nearly every neural area (**Fig. 1-7**). One system sends widespread projections from the pedunculopontine tegmentum and the laterodorsal pontine tegmentum to the thalamus and midbrain dopaminergic areas and also to the caudal pons and medulla (Woolf, 1991). The second cholinergic system arises from various basal forebrain nuclei projecting through the cerebral cortex and the

hippocampal formation (Woolf, 1991). These two systems provide broad and diffuse projections to wide areas of the brain. The third subsystem consists of striatal cholinergic interneurons which project throughout the striatum and the olfactory tubercle (Zhou et al., 2002).



**Fig. 1-7 Cholinergic circuits:** The basal forebrain contains two groups of cholinergic neurons: (1) the medial septal group (medial septal nucleus (**ms**) and vertical diagonal band (**vdb**)) that project cholinergic axons to the hippocampus and parahippocampal gyrus. (2) The nucleus basalis group (nucleus basalis (**bas**), substantia innominata (**si**) and horizontal diagonal band (**hdb**)) that project cholinergic axons to all parts of the neocortex, parts of limbic cortex and to the amygdala. The cholinergic pontomesencephalon neurons (laterodorsal tegmental (**ldt**) and pedunculopontine tegmental nuclei (**ppt**)) project onto hindbrain, thalamus, hypothalamus and basal forebrain. (Woolf, 1991)

In contrast to the main excitatory and inhibitory systems the cholinergic projections differ in their mode of transmitter release. While glutamate and GABA are released at synaptic sites, the majority of cortical and hippocampal cholinergic release sites are nonsynaptic (Umbriaco et al., 1994; Descarries et al., 1997). Nonsynaptic release and “spill over” of neurotransmitters from synapses contribute to volume transmission which enables ACh to diffuse and act at lower concentrations some distance away from the release site. The concentration of extrasynaptic ACh is regulated by AChE, which degrades ACh to choline and acetate. The presence of AChE ensures therefore that volume transmission is not uncontrolled, but rather that extrasynaptic ACh is cleared from distinct areas (Kawaja et al., 1990).

### 1.4.b. The $\alpha 7$ nAChR – unique among the nAChRs

Among the many possible compositions of nAChRs in the brain, the  $\alpha 7$  nAChR belongs to the most abundant variants, being just topped by the  $\alpha 4\beta 2$  nAChR. *In situ* hybridization studies detected high levels of  $\alpha 7$  nAChR transcripts in olfactory areas, hippocampus, amygdala, and cerebral cortex (Seguela et al., 1993). Nevertheless most of the initial studies on  $\alpha 7$  nAChR were done in the chick ciliary ganglion. The chick ciliary ganglion was favoured due to the size of the synapses, the high content of  $\alpha 7$  nAChRs, and the easy access to the tissue. There,  $\alpha 7$  nAChR is perisynaptically localized, excluded from the PSD (Jacob and Berg, 1983; Loring et al., 1985; Horch and Sargent, 1995).

Later studies focused on the hippocampal formation, where a fine network of cholinergic fibres innervates pyramidal cells, granule cells, interneurons, and neurons of the hilus (Frotscher and Leranth, 1985). But still, in addition to those direct synaptic contacts there is a high portion of volume transmission in the hippocampus (Umbriaco et al., 1994; Descarries et al., 1997).  $\alpha 7$  nAChRs were best investigated in the hippocampus where they are most prominent on GABAergic interneurons. In addition, the  $\alpha 7$  nAChR was found nearly on every synapse in the CA1 stratum radiatum of the rat hippocampus (Fabian-Fine et al., 2001). The largest size  $\alpha 7$  nAChR-containing interneurons are found in CA3, in both stratum radiatum and stratum lucidum (Freedman et al., 1993).  $\alpha 7$  nAChR differs in many aspects from other nAChRs. While the  $\alpha 4\beta 2$  nAChR has a high affinity for ACh but slow desensitization, the  $\alpha 7$  nAChR binds ACh with low affinity but has an exceptionally fast desensitization (Quick and Lester, 2002). These properties make  $\alpha 7$  nAChR a candidate responsive to volume transmission with low ACh levels featuring rare desensitization (Dani and Bertrand, 2007). Moreover  $\alpha 7$  nAChR is unique in terms of structure and permeability. To date,  $\alpha 7$  nAChR has been found to be the only homopentameric AChR in the brain with the highest affinity of all neuronal AChRs to  $\alpha$ -BT (Couturier et al., 1990; Seguela et al., 1993). Further, the  $\alpha 7$  nAChR differs strikingly from its homologues by displaying a much higher permeability of  $\text{Ca}^{2+}$  ions reaching values of the NMDAR in human brain

(Bertrand et al., 1993; Seguela et al., 1993). But unlike the NMDA-receptor, the  $\alpha 7$  nAChR is permeable for calcium ions even at resting membrane potential. These features of  $\alpha 7$  nAChR are highly conserved throughout the animal kingdom, pointing to a critical role of the receptor in neuronal networks (Le Novere et al., 2002).

#### **1.4.c. Nicotine, a modulator of cognitive function**

Tobacco smoking and acute and chronic administration of nicotine improve cognitive functions (Poincheval-Fuhrman and Sara, 1993). It has been shown that the improvement results from facilitated induction of long-term potentiation (LTP) in the hippocampus (Fujii et al., 1999), a synaptic model of learning and memory. The underlying mechanism is thought to be an ACh-dependent alteration of the oscillatory rhythms that result from the interaction of the various hippocampal subregions (Hasselmo, 2006). Altering such rhythms may result in an increase in the recruitment of various cell-signalling molecules known to be involved in synaptic plasticity as measured by LTP. Thereby different nAChR subtypes, mainly  $\alpha 7$  and  $\alpha 4\beta 2$  nAChRs, modulate different types of LTP in the various subregions of the hippocampus (Kenney and Gould, 2008).

While in rats injection of the  $\alpha 7$  nAChR antagonist MLA into the ventral hippocampus and basolateral amygdala impairs working memory (Levin, 2002), the administration of the partial  $\alpha 7$  nAChR agonist GTS-21 significantly improves learning and memory in rats (Arendash et al., 1995) and also attenuates the age-related impairment of classical conditioning in rabbits (Woodruff-Pak et al., 2000). A possible mechanism of nicotine induced LTP acts through the desensitization of  $\alpha 7$  nAChRs on inhibitory interneurons in the hippocampal CA1 by nicotine. The desensitization of  $\alpha 7$  nAChR reduces GABAergic inhibition and therefore indirectly increases the excitability of pyramidal cells, leading to LTP induction (Fujii et al., 2000).

Additional proof for the importance of  $\alpha 7$  nAChR in cognition was given with studies in  $\alpha 7$  nAChR deficient mice. In comparison to wild-type (WT) mice,  $\alpha 7$  nAChR knock-out (KO) mice showed significant deficits in attention and working memory. These behavioural deficits were associated with the loss of

$\alpha 7$  nAChR, as  $\alpha 4\beta 2$  nAChR density was unaltered in these mice (Young et al., 2007).

In addition to cognitive functions nicotine modulates also the dopaminergic (DA) system, which is involved in the reward circuits of the nucleus accumbens, an area involved in the development of addiction (Mansvelder and McGehee, 2002).  $\alpha 7$  nAChR KO mice were used to investigate the role of  $\alpha 7$  nAChR in nicotine addiction. Pons et al. showed that  $\alpha 7$  nAChR KO mice do self-administer nicotine in a manner indistinguishable from WT mice (Pons et al., 2008). This is consistent with the finding that  $\alpha 7$  nAChR is not involved in chronic nicotine self-administration (SA) in rats (Grottick et al., 2000).  $\alpha 7$  nAChR is believed to play a key role in long-term adaptations to passive chronic nicotine SA (Besson et al., 2007; Salas et al., 2007). Recent work performed in slices showed that following short applications of nicotine,  $\alpha 4\beta 2$  nAChRs on ventral tegmental area (VTA) DA neurons and VTA GABAergic neurons desensitize. However  $\alpha 7$  nAChRs on glutamatergic axons in the VTA do not desensitize and therefore potentiate glutamatergic excitation on DA neurons (Mansvelder and McGehee, 2000, 2002). As a consequence, DA neurons fire more frequently. The enhanced glutamatergic release produces LTP of the glutamatergic afferents contributing to the process of addiction (Dani and Harris, 2005). In addition to the existing evidence of the *in vitro* studies it is of note that a recent work showed that  $\alpha 7$  nAChR KO mice exhibit only a slightly altered electrophysiological response to nicotine in DAergic neurons *in vivo*.

Altogether  $\alpha 7$  nAChR involvement has been confirmed for cognitive functions such as attention and learning and memory. Although the implication of  $\alpha 7$  nAChR in nicotine addiction is controversial, several other diseases are linked to  $\alpha 7$  nAChR underlining its significance.

#### **1.4.d. Mental diseases and the involvement of $\alpha 7$ nAChR**

The improvement of cognitive functions due to nicotine administration was also perceived by patients suffering from schizophrenia. The prevalence of tobacco smokers in this group is significantly increased compared to healthy

subjects, pointing to a positive effect of nicotine attenuating the symptoms (de Leon et al., 1995). In addition to hallucinations and delusions, schizophrenia patients suffer from the ability to focus attention (Nuechterlein and Dawson, 1984). The latter symptom is regulated by  $\alpha 7$  nAChR. Pharmacological and genetical studies identified the  $\alpha 7$  nAChR as an important target in the treatment of schizophrenia (Freedman et al., 1994; Leonard et al., 2002). Patients with schizophrenia have altered expression and regulation of  $\alpha 7$  nAChRs in the reticular nucleus of the thalamus, the hippocampus, the cingulate cortex and the frontal lobe regions (Freedman et al., 1995; Court et al., 1999; Guan et al., 1999; Marutle et al., 2001). The underlying mechanism in attention deficiencies in schizophrenic patients involves  $\alpha 7$  nAChRs on GABAergic interneurons (Albuquerque et al., 1998; Frazier et al., 1998b).  $\alpha 7$  nAChR induces GABA release which acts on GABA<sub>B</sub>Rs thus decreasing the release of glutamate. This prevents hippocampal neurons acting properly in attention related tasks (Hershman et al., 1995). In addition to schizophrenia, the  $\alpha 7$  nAChR has been implicated in a wide range of neuronal dysfunction and mental illness (Picciotto and Zoli, 2002; Gotti and Clementi, 2004).

Alzheimer's disease (AD) is the most common form of degenerative dementia affecting mainly elderly persons (Palmer, 2002). In AD patients a progressive loss of cognitive functions particularly in learning and memory is correlated with the neuronal loss in the cholinergic system (Coyle et al., 1983; Perry, 1986). The most dramatic symptoms of AD appear with the loss of cholinergic neurons in the cortex and the hippocampus (Nordberg, 1999; Perry et al., 2000). To date no cure is known for AD, but treatment of AD patients with AChE inhibitors counteract against the loss of cholinergic signalling by decreasing the hydrolysis rate of ACh (Maelicke et al., 2001). The treatment with nicotinic agents improves cognitive deficits of AD patients (Nordberg, 1999; Picciotto and Zoli, 2002). Amongst all AChRs the  $\alpha 7$  nAChR became the main candidate playing a potential role in AD.  $\alpha 7$  nAChR has been found to bind with high affinity to the A $\beta_{1-42}$  (Wang et al., 2000; Oddo and LaFerla, 2006). Further support for this prospect was given by the discovery of  $\alpha 7$  nAChRs localized in amyloid plaques (Wang et al., 2000). There is accumulating evidence that A $\beta_{1-42}$  inhibits currents mediated by  $\alpha 7$  nAChRs

(Liu et al., 2001a; Pettit et al., 2001). The inhibition by A $\beta$  would explain the decreasing synaptic induction of LTP contributing to a cognitive decline (Freir et al., 2001). Additionally, transgenic AD mice displayed a loss of  $\alpha 7$  nAChRs (Oddo and LaFerla, 2006). All these findings bear evidence that  $\alpha 7$  nAChRs are directly involved in the cognitive decline during AD.

#### **1.4.e. Where is $\alpha 7$ nAChR?**

To further identify and understand the mechanisms of  $\alpha 7$  nAChR involvement in neuronal diseases and LTP induction, the knowledge of the receptor's cellular localization and the molecular interaction partners is crucial.

Up to date,  $\alpha 7$  nAChRs have been identified in postsynaptic, presynaptic, perisynaptic, and extrasynaptic sites. In the hippocampus and neocortex, notably on GABAergic interneurons (Freedman et al., 1993),  $\alpha 7$  nAChRs mediate cholinergic postsynaptic input (Frazier et al., 1998a) and enhance GABAergic IPSCs in principal neurons (Frazier et al., 2003). Ultrastructural studies reported their presence in glutamatergic postsynaptic sites on neurons of the somatosensory cortex (Levy and Aoki, 2002; Jones et al., 2004) but also in a perisynaptic annulus around dendritic glutamatergic postsynaptic spines on cortical pyramidal cells (Fabian-Fine et al., 2001). Possible functions of postsynaptic  $\alpha 7$  nAChR are assistance of NMDAR activation and LTP induction by removal of the Mg<sup>2+</sup> block, a modulatory role by Ca<sup>2+</sup> influx activating 2<sup>nd</sup> messenger pathways, and less likely depolarization of the membrane by  $\alpha 7$  nAChR itself. Besides postsynaptic  $\alpha 7$  nAChR, the receptor is also located presynaptically. There it facilitates vesicle fusion and the release of various neurotransmitters (McGehee et al., 1995; Alkondon et al., 1997; Li et al., 1998; Radcliffe and Dani, 1998; Maggi et al., 2001; Jones et al., 2004). In primary hippocampal cultures,  $\alpha 7$  nAChRs are prominent in interneurons, forming somato-dendritic clusters, partially localized at GABAergic synapses (Kawai et al., 2002). Recently, several lines of evidence indicated that  $\alpha 7$  nAChRs are expressed extrasynaptically in the CNS (Fabian-Fine et al., 2001; Jones and Wonnacott, 2004). Extrasynaptic  $\alpha 7$  nAChR may be targeted by ACh released into the extrasynaptic space. It has been shown that activation of extrasynaptic  $\alpha 7$  nAChRs by cholinergic

agonists either facilitates or depresses backpropagating action potentials, depending on the timing of  $\alpha 7$  nAChR activation (Rozsa et al., 2008). To localize  $\alpha 7$  nAChR at various sites, mechanisms are needed to target the receptor specifically. Adaptor proteins interacting with receptors are known to fulfil such a task. E.g. Gephyrin is the scaffolding protein of GABA<sub>A</sub>Rs stabilizing the receptor in inhibitory postsynapses (Essrich et al., 1998).

No  $\alpha 7$  nAChR interacting proteins apart from RIC-3 and Src-family kinases (SFKs) have been identified so far. However RIC-3 is thought to be responsible for the correct folding of  $\alpha 7$  nAChR. Therefore RIC-3 is probably not involved in membrane localization of  $\alpha 7$  nAChR (Millar, 2008). SFKs have been shown to phosphorylate  $\alpha 7$  nAChRs, causing decreased receptor activity (Charpantier et al., 2005). So far interaction partners regulating localization and clustering of  $\alpha 7$  nAChR are unknown.

Taken together,  $\alpha 7$  nAChR has been found in synaptic and non-synaptic sites. The evidence for presynaptic  $\alpha 7$  nAChR is based on many studies using different approaches, whereas the postsynaptic localization of  $\alpha 7$  nAChRs is still discussed.  $\alpha 7$  nAChRs identified in postsynapses by immunogold labeling are located certainly in noncholinergic synapses, presumably GABAergic or glutamatergic, because they include nearly all of the recognizable synapses in the CA1 region (Fabian-Fine et al., 2001). Furthermore,  $\alpha 7$  nAChRs cluster at GABAergic synapses on cultured hippocampal neurons in postsynaptic location (Kawai et al., 2002). Despite all the published results showing  $\alpha 7$  nAChR postsynaptically localized, the receptor may be viewed as perisynaptic where it modulates effective noncholinergic synapses (Berg and Conroy, 2002).



## 1.5. Aim

In this thesis we try to uncover the mechanisms regulating the membrane dynamics and the clustering of  $\alpha 7$  nAChRs.

The clustering of a membrane protein is dependent on the stabilization by a direct or indirect link to scaffolding elements. The indirect link is performed by adaptor proteins binding to the membrane protein and scaffolding protein, thus forming a stable complex.

In chapter 2 we hypothesize the existence and function of a potential  $\alpha 7$  nAChR interacting protein. Does this protein regulate  $\alpha 7$  nAChR clustering and distribution? Which domains are essential for protein-protein binding? To address these questions we first identified  $\alpha 7$  nAChR interacting proteins by a yeast two-hybrid assay, using the large cytoplasmatic loop of  $\alpha 7$  nAChR as bait. PICK1, a protein known to interact with  $\text{PKC}\alpha$ , was identified as  $\alpha 7$  nAChR interacting protein. Immunoprecipitation (IP) experiments were performed in the rat brain and cell lines to investigate the protein-protein interaction and the involved protein domains. To focus on the functional role of PICK1 on  $\alpha 7$  nAChR, PICK1 overexpression in rat hippocampal neurons using virus and lipofection techniques was carried out.

The regulation of receptor clustering and localization is consequently also influenced by the membrane diffusion of a receptor. Membrane proteins have been shown to diffuse within the membrane of the cell surface. Areas where diffusion is decreased are spots where membrane proteins aggregate. To understand the regulation of  $\alpha 7$  nAChR clustering, it is not only important to know the interacting proteins but also to investigate the dynamics of the receptor within the membrane. To enable such an investigation, it is necessary to express different exogenous-tagged proteins in specific neurons. Transfected gene constructs can code for fluorescently-tagged proteins or for mutant proteins, used as markers or tools to interfere with cellular mechanisms, respectively. At the time I started my thesis work, there was no satisfying method known to transiently transfect cultured neurons with high efficiency and low toxicity. In chapter 3 we describe the development and optimization of a protocol for the efficient transfection of cultured neurons.

Having achieved the capability to transiently transfect neurons the investigation of surface  $\alpha 7$  nAChR dynamics became possible. In chapter 4 we focused on  $\alpha 7$  nAChRs in living neurons. We tackled the questions how mobile  $\alpha 7$  nAChRs are in the cell membrane? Is the mobility affected by scaffolding elements such as microtubules or the actin network? At which sites is  $\alpha 7$  nAChR mobility decreased? Is the mobility of  $\alpha 7$  nAChR affected by neuronal activity? To answer these questions single  $\alpha 7$  nAChRs were labeled with QDOTs, which are small fluorescent probes with a high quantum yield and high photostability, and tracked by live microscopy. Inhibitory and excitatory PSDs were visualized by the expression of fluorescently tagged marker proteins in the  $\alpha 7$  nAChR positive neurons. Receptor mobility in correlation to their localization was analyzed by custom software.

In this thesis work we try to reveal the mechanisms of  $\alpha 7$  nAChR clustering, possible interacting proteins, the underlying dynamic diffusion behavior of single  $\alpha 7$  nAChR receptors, and their adaption to neuronal activity. The answers to these questions will hopefully help to clarify the ongoing debate on  $\alpha 7$  nAChR and allow new speculations and hypotheses.

## References

- Albuquerque EX, Pereira EF, Braga MF, Alkondon M (1998) Contribution of nicotinic receptors to the function of synapses in the central nervous system: the action of choline as a selective agonist of alpha 7 receptors. *J Physiol Paris* 92:309-316.
- Alkondon M, Pereira EF, Albuquerque EX (1998) alpha-bungarotoxin- and methyllycaconitine-sensitive nicotinic receptors mediate fast synaptic transmission in interneurons of rat hippocampal slices. *Brain Res* 810:257-263.
- Alkondon M, Pereira EF, Barbosa CT, Albuquerque EX (1997) Neuronal nicotinic acetylcholine receptor activation modulates gamma-aminobutyric acid release from CA1 neurons of rat hippocampal slices. *J Pharmacol Exp Ther* 283:1396-1411.
- Allison DW, Gelfand VI, Spector I, Craig AM (1998) Role of actin in anchoring postsynaptic receptors in cultured hippocampal neurons: differential attachment of NMDA versus AMPA receptors. *J Neurosci* 18:2423-2436.
- Archibald K, Perry MJ, Molnar E, Henley JM (1998) Surface expression and metabolic half-life of AMPA receptors in cultured rat cerebellar granule cells. *Neuropharmacology* 37:1345-1353.
- Arendash GW, Sengstock GJ, Sanberg PR, Kem WR (1995) Improved learning and memory in aged rats with chronic administration of the nicotinic receptor agonist GTS-21. *Brain Res* 674:252-259.
- Baer K, Burli T, Huh KH, Wiesner A, Erb-Vogtli S, Gockeritz-Dujmovic D, Moransard M, Nishimune A, Rees MI, Henley JM, Fritschy JM, Fuhrer C (2007) PICK1 interacts with alpha7 neuronal nicotinic acetylcholine receptors and controls their clustering. *Mol Cell Neurosci* 35:339-355.
- Bannai H, Levi S, Schweizer C, Dahan M, Triller A (2006) Imaging the lateral diffusion of membrane molecules with quantum dots. *Nat Protoc* 1:2628-2634.
- Bats C, Choquet D (2007) The interaction between Stargazin and PSD-95 regulates AMPA receptor surface trafficking. *Neuron* 53:719-734.
- Ben-Ami HC, Yassin L, Farah H, Michaeli A, Eshel M, Treinin M (2005) RIC-3 affects properties and quantity of nicotinic acetylcholine receptors via a mechanism that does not require the coiled-coil domains. *J Biol Chem* 280:28053-28060.
- Bennett V, Gilligan DM (1993) The spectrin-based membrane skeleton and micron-scale organization of the plasma membrane. *Annu Rev Cell Biol* 9:27-66.
- Berg DK, Conroy WG (2002) Nicotinic alpha 7 receptors: synaptic options and downstream signaling in neurons. *J Neurobiol* 53:512-523.
- Bernard C (1850) Recherches sur le curare. Written with J. Pelouze. *Comptes rendus hebdomadaires de l'Académie des sciences, Paris* 1850, 31: 533-537.
- Bertrand D, Galzi JL, Devillers-Thiery A, Bertrand S, Changeux JP (1993) Mutations at two distinct sites within the channel domain M2 alter calcium permeability of neuronal alpha 7 nicotinic receptor. *Proc Natl Acad Sci U S A* 90:6971-6975.
- Besson M, Granon S, Mameli-Engvall M, Cloez-Tayarani I, Maubourguet N, Cormier A, Cazala P, David V, Changeux JP, Faure P (2007) Long-term effects of chronic nicotine exposure on brain nicotinic receptors. *Proc Natl Acad Sci U S A* 104:8155-8160.
- Blanpied TA, Scott DB, Ehlers MD (2002) Dynamics and regulation of clathrin coats at specialized endocytic zones of dendrites and spines. *Neuron* 36:435-449.
- Boudin H, Craig AM (2001) Molecular determinants for PICK1 synaptic aggregation and mGluR7a receptor coclustering: role of the PDZ, coiled-coil, and acidic domains. *J Biol Chem* 276:30270-30276.
- Boudin H, Doan A, Xia J, Shigemoto R, Huganir RL, Worley P, Craig AM (2000) Presynaptic clustering of mGluR7a requires the PICK1 PDZ domain binding site. *Neuron* 28:485-497.
- Brunig I, Scotti E, Sidler C, Fritschy JM (2002a) Intact sorting, targeting, and clustering of gamma-aminobutyric acid A receptor subtypes in hippocampal neurons in vitro. *J Comp Neurol* 443:43-55.
- Brunig I, Suter A, Knuesel I, Luscher B, Fritschy JM (2002b) GABAergic terminals are required for postsynaptic clustering of dystrophin but not of GABA(A) receptors and gephyrin. *J Neurosci* 22:4805-4813.
- Bruses JL, Chauvet N, Rutishauser U (2001) Membrane lipid rafts are necessary for the maintenance of the (alpha)7 nicotinic acetylcholine receptor in somatic spines of ciliary neurons. *J Neurosci* 21:504-512.
- Buerli T, Pellegrino C, Baer K, Lardi-Studler B, Chudotvorova I, Fritschy JM, Medina I, Fuhrer C (2007) Efficient transfection of DNA or shRNA vectors into neurons using magnetofection. *Nat Protoc* 2:3090-3101.
- Carlin RK, Grab DJ, Cohen RS, Siekevitz P (1980) Isolation and characterization of postsynaptic densities from various brain regions: enrichment of different types of postsynaptic densities. *J Cell Biol* 86:831-845.
- Chalfie M, Tu Y, Euskirchen G, Ward WW, Prasher DC (1994) Green fluorescent protein as a marker for gene expression. *Science* 263:802-805.
- Chan WC, Nie S (1998) Quantum dot bioconjugates for ultrasensitive nonisotopic detection. *Science* 281:2016-2018.
- Chang CC, Lee CY (1963) Isolation of Neurotoxins from the Venom of *Bungarus Multicinctus* and Their Modes of Neuromuscular Blocking Action. *Arch Int Pharmacodyn Ther* 144:241-257.
- Chang KT, Berg DK (2001) Voltage-gated channels block nicotinic regulation of CREB phosphorylation and gene expression in neurons. *Neuron* 32:855-865.
- Changeux JP, Kasai M, Lee CY (1970) Use of a snake venom toxin to characterize the cholinergic receptor protein. *Proc Natl Acad Sci U S A* 67:1241-1247.
- Charpentier E, Wiesner A, Huh KH, Ogier R, Hoda JC, Allaman G, Raggenbass M, Feuerbach D, Bertrand D, Fuhrer C (2005) Alpha7 neuronal nicotinic acetylcholine receptors are negatively regulated by tyrosine phosphorylation and Src-family kinases. *J Neurosci* 25:9836-9849.
- Charrier C, Ehrensperger MV, Dahan M, Levi S, Triller A (2006) Cytoskeleton regulation of glycine receptor number at synapses and diffusion in the plasma membrane. *J Neurosci* 26:8502-8511.
- Chen D, Patrick JW (1997) The alpha-bungarotoxin-binding nicotinic acetylcholine receptor from rat brain contains only the alpha7 subunit. *J Biol Chem* 272:24024-24029.
- Chen HJ, Rojas-Soto M, Oguni A, Kennedy MB (1998) A synaptic Ras-GTPase activating protein (p135 SynGAP) inhibited by CaM kinase II. *Neuron* 20:895-904.

- Cho CH, Song W, Leitzell K, Teo E, Meleth AD, Quick MW, Lester RA (2005) Rapid upregulation of  $\alpha 7$  nicotinic acetylcholine receptors by tyrosine dephosphorylation. *J Neurosci* 25:3712-3723.
- Cho KO, Hunt CA, Kennedy MB (1992) The rat brain postsynaptic density fraction contains a homolog of the *Drosophila* discs-large tumor suppressor protein. *Neuron* 9:929-942.
- Chudotvorova I, Ivanov A, Rama S, Hubner CA, Pellegrino C, Ben-Ari Y, Medina I (2005) Early expression of KCC2 in rat hippocampal cultures augments expression of functional GABA synapses. *J Physiol* 566:671-679.
- Cincotta SL, Yorek MS, Moschak TM, Lewis SR, Rodefer JS (2008) Selective nicotinic acetylcholine receptor agonists: potential therapies for neuropsychiatric disorders with cognitive dysfunction. *Curr Opin Investig Drugs* 9:47-56.
- Claudio T, Ballivet M, Patrick J, Heinemann S (1983) Nucleotide and deduced amino acid sequences of Torpedo californica acetylcholine receptor gamma subunit. *Proc Natl Acad Sci U S A* 80:1111-1115.
- Conroy WG, Liu Z, Nai Q, Coggan JS, Berg DK (2003) PDZ-containing proteins provide a functional postsynaptic scaffold for nicotinic receptors in neurons. *Neuron* 38:759-771.
- Cooper ST, Millar NS (1997) Host cell-specific folding and assembly of the neuronal nicotinic acetylcholine receptor  $\alpha 7$  subunit. *J Neurochem* 68:2140-2151.
- Court J, Spurdin D, Lloyd S, McKeith I, Ballard C, Cairns N, Kerwin R, Perry R, Perry E (1999) Neuronal nicotinic receptors in dementia with Lewy bodies and schizophrenia: alpha-bungarotoxin and nicotine binding in the thalamus. *J Neurochem* 73:1590-1597.
- Couturier S, Bertrand D, Matter JM, Hernandez MC, Bertrand S, Millar N, Valera S, Barkas T, Ballivet M (1990) A neuronal nicotinic acetylcholine receptor subunit ( $\alpha 7$ ) is developmentally regulated and forms a homo-oligomeric channel blocked by alpha-BTX. *Neuron* 5:847-856.
- Cowan CA, Yokoyama N, Bianchi LM, Henkemeyer M, Fritsch B (2000) EphB2 guides axons at the midline and is necessary for normal vestibular function. *Neuron* 26:417-430.
- Coyle JT, Price DL, DeLong MR (1983) Alzheimer's disease: a disorder of cortical cholinergic innervation. *Science* 219:1184-1190.
- Dahan M, Levi S, Luccardini C, Rostaing P, Riveau B, Triller A (2003) Diffusion dynamics of glycine receptors revealed by single-quantum dot tracking. *Science* 302:442-445.
- Dajas-Bailador F, Wonnacott S (2004) Nicotinic acetylcholine receptors and the regulation of neuronal signalling. *Trends Pharmacol Sci* 25:317-324.
- Dale H (1914b) The occurrence in ergot and action of acetyl-choline. *Journal of Physiology* 48:3-4.
- Dani JA, Harris RA (2005) Nicotine addiction and comorbidity with alcohol abuse and mental illness. *Nat Neurosci* 8:1465-1470.
- Dani JA, Bertrand D (2007) Nicotinic acetylcholine receptors and nicotinic cholinergic mechanisms of the central nervous system. *Annu Rev Pharmacol Toxicol* 47:699-729.
- de Leon J, Dadvand M, Canuso C, White AO, Stanilla JK, Simpson GM (1995) Schizophrenia and smoking: an epidemiological survey in a state hospital. *Am J Psychiatry* 152:453-455.
- Debanne D, Daoudal G, Sourdet V, Russier M (2003) Brain plasticity and ion channels. *J Physiol Paris* 97:403-414.
- Descarries L, Gisiger V, Steriade M (1997) Diffuse transmission by acetylcholine in the CNS. *Prog Neurobiol* 53:603-625.
- Dev KK, Nishimune A, Henley JM, Nakanishi S (1999) The protein kinase C  $\alpha$  binding protein PICK1 interacts with short but not long form alternative splice variants of AMPA receptor subunits. *Neuropharmacology* 38:635-644.
- Dev KK, Nakajima Y, Kitano J, Braithwaite SP, Henley JM, Nakanishi S (2000) PICK1 interacts with and regulates PKC phosphorylation of mGluR7. *J Neurosci* 20:7252-7257.
- Dichter MA, Ayala GF (1987) Cellular mechanisms of epilepsy: a status report. *Science* 237:157-164.
- Drisdel RC, Green WN (2000) Neuronal alpha-bungarotoxin receptors are  $\alpha 7$  subunit homomers. *J Neurosci* 20:133-139.
- Drisdel RC, Manzana E, Green WN (2004) The role of palmitoylation in functional expression of nicotinic  $\alpha 7$  receptors. *J Neurosci* 24:10502-10510.
- Duclos S, Diez R, Garin J, Papadopoulou B, Descoteaux A, Stenmark H, Desjardins M (2000) Rab5 regulates the kiss and run fusion between phagosomes and endosomes and the acquisition of phagosome leishmanicidal properties in RAW 264.7 macrophages. *J Cell Sci* 113 Pt 19:3531-3541.
- Dunaevsky A, Tashiro A, Majewska A, Mason C, Yuste R (1999) Developmental regulation of spine motility in the mammalian central nervous system. *Proc Natl Acad Sci U S A* 96:13438-13443.
- Eglen RM (2006) Muscarinic receptor subtypes in neuronal and non-neuronal cholinergic function. *Auton Autacoid Pharmacol* 26:219-233.
- Ehlers MD, Heine M, Groc L, Lee MC, Choquet D (2007) Diffusional trapping of GluR1 AMPA receptors by input-specific synaptic activity. *Neuron* 54:447-460.
- Ehrensperger MV, Hanus C, Vannier C, Triller A, Dahan M (2007) Multiple association states between glycine receptors and gephyrin identified by SPT analysis. *Biophys J* 92:3706-3718.
- Eitzen G (2003) Actin remodeling to facilitate membrane fusion. *Biochim Biophys Acta* 1641:175-181.
- Essrich C, Lorez M, Benson JA, Fritschy JM, Luscher B (1998) Postsynaptic clustering of major GABAA receptor subtypes requires the gamma 2 subunit and gephyrin. *Nat Neurosci* 1:563-571.
- Fabian-Fine R, Skehel P, Errington ML, Davies HA, Sher E, Stewart MG, Fine A (2001) Ultrastructural distribution of the  $\alpha 7$  nicotinic acetylcholine receptor subunit in rat hippocampus. *J Neurosci* 21:7993-8003.
- Fields S, Song O (1989) A novel genetic system to detect protein-protein interactions. *Nature* 340:245-246.
- Frazier CJ, Strowbridge BW, Papke RL (2003) Nicotinic receptors on local circuit neurons in dentate gyrus: a potential role in regulation of granule cell excitability. *J Neurophysiol* 89:3018-3028.
- Frazier CJ, Buhler AV, Weiner JL, Dunwiddie TV (1998a) Synaptic potentials mediated via alpha-bungarotoxin-sensitive nicotinic acetylcholine receptors in rat hippocampal interneurons. *J Neurosci* 18:8228-8235.
- Frazier CJ, Rollins YD, Breese CR, Leonard S, Freedman R, Dunwiddie TV (1998b) Acetylcholine activates an alpha-bungarotoxin-sensitive nicotinic current in rat hippocampal interneurons, but not pyramidal cells. *J Neurosci* 18:1187-1195.
- Freedman R, Hall M, Adler LE, Leonard S (1995) Evidence in postmortem brain tissue for decreased numbers of hippocampal nicotinic receptors in schizophrenia. *Biol Psychiatry* 38:22-33.

- Freedman R, Wetmore C, Stromberg I, Leonard S, Olson L (1993) Alpha-bungarotoxin binding to hippocampal interneurons: immunocytochemical characterization and effects on growth factor expression. *J Neurosci* 13:1965-1975.
- Freedman R, Adler LE, Bickford P, Byerley W, Coon H, Cullum CM, Griffith JM, Harris JG, Leonard S, Miller C, et al. (1994) Schizophrenia and nicotinic receptors. *Harv Rev Psychiatry* 2:179-192.
- Freir DB, Holscher C, Herron CE (2001) Blockade of long-term potentiation by beta-amyloid peptides in the CA1 region of the rat hippocampus in vivo. *J Neurophysiol* 85:708-713.
- Fritschy JM, Mohler H (1995) GABAA-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J Comp Neurol* 359:154-194.
- Frotscher M, Leranth C (1985) Cholinergic innervation of the rat hippocampus as revealed by choline acetyltransferase immunocytochemistry: a combined light and electron microscopic study. *J Comp Neurol* 239:237-246.
- Fuhrer C, Hall ZW (1996) Functional interaction of Src family kinases with the acetylcholine receptor in C2 myotubes. *J Biol Chem* 271:32474-32481.
- Fuhrer C, Sugiyama JE, Taylor RG, Hall ZW (1997) Association of muscle-specific kinase MuSK with the acetylcholine receptor in mammalian muscle. *Embo J* 16:4951-4960.
- Fujii K, Maeda K, Hikida T, Mustafa AK, Balkissoon R, Xia J, Yamada T, Ozeki Y, Kawahara R, Okawa M, Huganir RL, Ujike H, Snyder SH, Sawa A (2006) Serine racemase binds to PICK1: potential relevance to schizophrenia. *Mol Psychiatry* 11:150-157.
- Fujii S, Ji Z, Morita N, Sumikawa K (1999) Acute and chronic nicotine exposure differentially facilitate the induction of LTP. *Brain Res* 846:137-143.
- Fujii S, Jia Y, Yang A, Sumikawa K (2000) Nicotine reverses GABAergic inhibition of long-term potentiation induction in the hippocampal CA1 region. *Brain Res* 863:259-265.
- Gahring LC, Rogers SW (2005) Neuronal nicotinic acetylcholine receptor expression and function on nonneuronal cells. *Aaps J* 7:E885-894.
- Gaiarsa JL, Caillard O, Ben-Ari Y (2002) Long-term plasticity at GABAergic and glycinergic synapses: mechanisms and functional significance. *Trends Neurosci* 25:564-570.
- Gautam M, Noakes PG, Mudd J, Nichol M, Chu GC, Sanes JR, Merlie JP (1995) Failure of postsynaptic specialization to develop at neuromuscular junctions of rapsyn-deficient mice. *Nature* 377:232-236.
- Gee SH, Sekely SA, Lombardo C, Kurakin A, Froehner SC, Kay BK (1998) Cyclic peptides as non-carboxyl-terminal ligands of syntrophin PDZ domains. *J Biol Chem* 273:21980-21987.
- Gingras J, Rassadi S, Cooper E, Ferns M (2002) Agrin plays an organizing role in the formation of sympathetic synapses. *J Cell Biol* 158:1109-1118.
- Glendenning KK (1998) Thalamic inhibition in the evolution of human intelligence : Evolutionary pressure for cortical inhibition. *Mankind quarterly* 38:319-335 (312 p.311/312).
- Goetze B, Grunewald B, Baldassa S, Kiebler M (2004) Chemically controlled formation of a DNA/calcium phosphate coprecipitate: application for transfection of mature hippocampal neurons. *J Neurobiol* 60:517-525.
- Gotti C, Clementi F (2004) Neuronal nicotinic receptors: from structure to pathology. *Prog Neurobiol* 74:363-396.
- Groc L, Heine M, Cognet L, Brickley K, Stephenson FA, Lounis B, Choquet D (2004) Differential activity-dependent regulation of the lateral mobilities of AMPA and NMDA receptors. *Nat Neurosci* 7:695-696.
- Groc L, Lafourcade M, Heine M, Renner M, Racine V, Sibarita JB, Lounis B, Choquet D, Cognet L (2007) Surface trafficking of neurotransmitter receptor: comparison between single-molecule/quantum dot strategies. *J Neurosci* 27:12433-12437.
- Grottick AJ, Trube G, Corrigan WA, Huwyler J, Malherbe P, Wyler R, Higgins GA (2000) Evidence that nicotinic alpha(7) receptors are not involved in the hyperlocomotor and rewarding effects of nicotine. *J Pharmacol Exp Ther* 294:1112-1119.
- Guan ZZ, Zhang X, Blennow K, Nordberg A (1999) Decreased protein level of nicotinic receptor alpha7 subunit in the frontal cortex from schizophrenic brain. *Neuroreport* 10:1779-1782.
- Gulyas AI, Megias M, Emri Z, Freund TF (1999) Total number and ratio of excitatory and inhibitory synapses converging onto single interneurons of different types in the CA1 area of the rat hippocampus. *J Neurosci* 19:10082-10097.
- Halevi S, McKay J, Palfreyman M, Yassin L, Eshel M, Jorgensen E, Treinin M (2002) The C. elegans ric-3 gene is required for maturation of nicotinic acetylcholine receptors. *Embo J* 21:1012-1020.
- Hanley JG (2008) PICK1: a multi-talented modulator of AMPA receptor trafficking. *Pharmacol Ther* 118:152-160.
- Harris BZ, Lim WA (2001) Mechanism and role of PDZ domains in signaling complex assembly. *J Cell Sci* 114:3219-3231.
- Hasselmo ME (2006) The role of acetylcholine in learning and memory. *Curr Opin Neurobiol* 16:710-715.
- Hebb DO (1949) *The Organization of Behavior: A Neuropsychological Theory* Wiley.
- Heine M, Groc L, Frischknecht R, Beique JC, Lounis B, Rumbaugh G, Huganir RL, Cognet L, Choquet D (2008) Surface mobility of postsynaptic AMPARs tunes synaptic transmission. *Science* 320:201-205.
- Hershman KM, Freedman R, Bickford PC (1995) GABAB antagonists diminish the inhibitory gating of auditory response in the rat hippocampus. *Neurosci Lett* 190:133-136.
- Hirbec H, Francis JC, Lauri SE, Braithwaite SP, Coussen F, Mulle C, Dev KK, Coutinho V, Meyer G, Isaac JT, Collingridge GL, Henley JM (2003) Rapid and differential regulation of AMPA and kainate receptors at hippocampal mossy fibre synapses by PICK1 and GRIP. *Neuron* 37:625-638.
- Hong CJ, Liao DL, Shih HL, Tsai SJ (2004) Association study of PICK1 rs3952 polymorphism and schizophrenia. *Neuroreport* 15:1965-1967.
- Horch HL, Sargent PB (1995) Perisynaptic surface distribution of multiple classes of nicotinic acetylcholine receptors on neurons in the chicken ciliary ganglion. *J Neurosci* 15:7778-7795.
- Hucho F, Changeux JP (1973) Molecular weight and quaternary structure of the cholinergic receptor protein extracted by detergents from Electrophorus electricus electric tissue. *FEBS Lett* 38:11-15.
- Huh KH, Fuhrer C (2002) Clustering of nicotinic acetylcholine receptors: from the neuromuscular junction to interneuronal synapses. *Mol Neurobiol* 25:79-112.
- Jacob MH, Berg DK (1983) The ultrastructural localization of alpha-bungarotoxin binding sites in relation to synapses on chick ciliary ganglion neurons. *J Neurosci* 3:260-271.

- Jacob TC, Moss SJ, Jurd R (2008) GABA(A) receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat Rev Neurosci* 9:331-343.
- Jacquier A, Buhler E, Schafer MK, Bohl D, Blanchard S, Beclin C, Haase G (2006) Alsln/Rac1 signaling controls survival and growth of spinal motoneurons. *Ann Neurol* 60:105-117.
- Ji D, Lape R, Dani JA (2001) Timing and location of nicotinic activity enhances or depresses hippocampal synaptic plasticity. *Neuron* 31:131-141.
- Jiang M, Chen G (2006) High Ca<sup>2+</sup>-phosphate transfection efficiency in low-density neuronal cultures. *Nat Protoc* 1:695-700.
- Jin W, Ge WP, Xu J, Cao M, Peng L, Yung W, Liao D, Duan S, Zhang M, Xia J (2006) Lipid binding regulates synaptic targeting of PICK1, AMPA receptor trafficking, and synaptic plasticity. *J Neurosci* 26:2380-2390.
- Jonas P, Burnashev N (1995) Molecular mechanisms controlling calcium entry through AMPA-type glutamate receptor channels. *Neuron* 15:987-990.
- Jones IW, Wonnacott S (2004) Precise localization of alpha7 nicotinic acetylcholine receptors on glutamatergic axon terminals in the rat ventral tegmental area. *J Neurosci* 24:11244-11252.
- Jones IW, Barik J, O'Neill MJ, Wonnacott S (2004) Alpha bungarotoxin-1.4 nm gold: a novel conjugate for visualising the precise subcellular distribution of alpha 7\* nicotinic acetylcholine receptors. *J Neurosci Methods* 134:65-74.
- Jones S, Yakel JL (1997) Functional nicotinic ACh receptors on interneurons in the rat hippocampus. *J Physiol* 504 (Pt 3):603-610.
- Jones S, Sudweeks S, Yakel JL (1999) Nicotinic receptors in the brain: correlating physiology with function. *Trends Neurosci* 22:555-561.
- Kandel ER, Schwartz JH, Jessell TM (2000) Principles of neural science. 4th edition.
- Karlin A, Cox RN, Dipaola M, Holtzman E, Kao PN, Lobel P, Wang L, Yodh N (1986) Functional domains of the nicotinic acetylcholine receptor. *Ann N Y Acad Sci* 463:53-69.
- Kawaguchi SY, Hirano T (2002) Signaling cascade regulating long-term potentiation of GABA(A) receptor responsiveness in cerebellar Purkinje neurons. *J Neurosci* 22:3969-3976.
- Kawai H, Zago W, Berg DK (2002) Nicotinic alpha 7 receptor clusters on hippocampal GABAergic neurons: regulation by synaptic activity and neurotrophins. *J Neurosci* 22:7903-7912.
- Kawaja MD, Flumerfelt BA, Hryciyshyn AW (1990) A comparison of the subnuclear and ultrastructural distribution of acetylcholinesterase and choline acetyltransferase in the rat interpeduncular nucleus. *Brain Res Bull* 24:517-523.
- Kawashima K, Fujii T (2003) The lymphocytic cholinergic system and its contribution to the regulation of immune activity. *Life Sci* 74:675-696.
- Kenney JW, Gould TJ (2008) Modulation of hippocampus-dependent learning and synaptic plasticity by nicotine. *Mol Neurobiol* 38:101-121.
- Kessler RM, Woodward ND, Riccardi P, Li R, Ansari MS, Anderson S, Dawant B, Zald D, Meltzer HY (2009) Dopamine D(2) Receptor Levels in Striatum, Thalamus, Substantia Nigra, Limbic Regions, and Cortex in Schizophrenic Subjects. *Biol Psychiatry*.
- Kim JB, Choi JS, Nam K, Lee M, Park JS, Lee JK (2006) Enhanced transfection of primary cortical cultures using arginine-grafted PAMAM dendrimer, PAMAM-Arg. *J Control Release* 114:110-117.
- Kim JH, Liao D, Lau LF, Haganir RL (1998) SynGAP: a synaptic RasGAP that associates with the PSD-95/SAP90 protein family. *Neuron* 20:683-691.
- Kim MJ, Dunah AW, Wang YT, Sheng M (2005) Differential roles of NR2A- and NR2B-containing NMDA receptors in Ras-ERK signaling and AMPA receptor trafficking. *Neuron* 46:745-760.
- Kneussel M, Betz H (2000) Receptors, gephyrin and gephyrin-associated proteins: novel insights into the assembly of inhibitory postsynaptic membrane specializations. *J Physiol* 525 Pt 1:1-9.
- Langley JN (1905) On the reaction of cells and of nerve-endings to certain poisons, chiefly as regards the reaction of striated muscle to nicotine and to curari. *J Physiol* 33:374-413.
- Lansdell SJ, Gee VJ, Harkness PC, Doward AI, Baker ER, Gibb AJ, Millar NS (2005) RIC-3 enhances functional expression of multiple nicotinic acetylcholine receptor subtypes in mammalian cells. *Mol Pharmacol* 68:1431-1438.
- Lardi-Studler B, Smolinsky B, Petitjean CM, Koenig F, Sidler C, Meier JC, Fritschy JM, Schwarz G (2007) Vertebrate-specific sequences in the gephyrin E-domain regulate cytosolic aggregation and postsynaptic clustering. *J Cell Sci* 120:1371-1382.
- Le Novère N, Corringer PJ, Changeux JP (2002) The diversity of subunit composition in nAChRs: evolutionary origins, physiologic and pharmacologic consequences. *J Neurobiol* 53:447-456.
- Leonard S, Gault J, Hopkins J, Logel J, Vianzon R, Short M, Drebing C, Berger R, Venn D, Sirota P, Zerbe G, Olincy A, Ross RG, Adler LE, Freedman R (2002) Association of promoter variants in the alpha7 nicotinic acetylcholine receptor subunit gene with an inhibitory deficit found in schizophrenia. *Arch Gen Psychiatry* 59:1085-1096.
- Leranth C, Szeideemann Z, Hsu M, Buzsaki G (1996) AMPA receptors in the rat and primate hippocampus: a possible absence of GluR2/3 subunits in most interneurons. *Neuroscience* 70:631-652.
- Levi S, Schweizer C, Bannai H, Pascual O, Charrier C, Triller A (2008) Homeostatic regulation of synaptic GlyR numbers driven by lateral diffusion. *Neuron* 59:261-273.
- Levin ED (2002) Nicotinic receptor subtypes and cognitive function. *J Neurobiol* 53:633-640.
- Levin ED, McClernon FJ, Rezvani AH (2006) Nicotinic effects on cognitive function: behavioral characterization, pharmacological specification, and anatomic localization. *Psychopharmacology (Berl)* 184:523-539.
- Levy RB, Aoki C (2002) Alpha7 nicotinic acetylcholine receptors occur at postsynaptic densities of AMPA receptor-positive and -negative excitatory synapses in rat sensory cortex. *J Neurosci* 22:5001-5015.
- Li X, Rainnie DG, McCarley RW, Greene RW (1998) Presynaptic nicotinic receptors facilitate monoaminergic transmission. *J Neurosci* 18:1904-1912.
- Lindstrom J, Anand R, Peng X, Gerzanich V, Wang F, Li Y (1995) Neuronal nicotinic receptor subtypes. *Ann N Y Acad Sci* 757:100-116.
- Liu Q, Kawai H, Berg DK (2001a) beta -Amyloid peptide blocks the response of alpha 7-containing nicotinic receptors on hippocampal neurons. *Proc Natl Acad Sci U S A* 98:4734-4739.

- Liu Y, Ford B, Mann MA, Fischbach GD (2001b) Neuregulins increase  $\alpha 7$  nicotinic acetylcholine receptors and enhance excitatory synaptic transmission in GABAergic interneurons of the hippocampus. *J Neurosci* 21:5660-5669.
- Loring RH, Zigmond RE (1987) Ultrastructural distribution of 125I-toxin F binding sites on chick ciliary neurons: synaptic localization of a toxin that blocks ganglionic nicotinic receptors. *J Neurosci* 7:2153-2162.
- Loring RH, Dahm LM, Zigmond RE (1985) Localization of alpha-bungarotoxin binding sites in the ciliary ganglion of the embryonic chick: an autoradiographic study at the light and electron microscopic level. *Neuroscience* 14:645-660.
- Lu J, Helton TD, Blanpied TA, Raczy B, Newpher TM, Weinberg RJ, Ehlers MD (2007) Postsynaptic positioning of endocytic zones and AMPA receptor cycling by physical coupling of dynamin-3 to Homer. *Neuron* 55:874-889.
- Madsen KL, Beuming T, Niv MY, Chang CW, Dev KK, Weinstein H, Gether U (2005) Molecular determinants for the complex binding specificity of the PDZ domain in PICK1. *J Biol Chem* 280:20539-20548.
- Maelicke A, Samochocki M, Jostock R, Fehrenbacher A, Ludwig J, Albuquerque EX, Zerlin M (2001) Allosteric sensitization of nicotinic receptors by galantamine, a new treatment strategy for Alzheimer's disease. *Biol Psychiatry* 49:279-288.
- Maggi L, Sher E, Cherubini E (2001) Regulation of GABA release by nicotinic acetylcholine receptors in the neonatal rat hippocampus. *J Physiol* 536:89-100.
- Maletic-Savatic M, Malinow R (1998) Calcium-evoked dendritic exocytosis in cultured hippocampal neurons. Part I: trans-Golgi network-derived organelles undergo regulated exocytosis. *J Neurosci* 18:6803-6813.
- Mansvelder HD, McGehee DS (2000) Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* 27:349-357.
- Mansvelder HD, McGehee DS (2002) Cellular and synaptic mechanisms of nicotine addiction. *J Neurobiol* 53:606-617.
- Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C (2004) Interneurons of the neocortical inhibitory system. *Nat Rev Neurosci* 5:793-807.
- Martin LF, Kem WR, Freedman R (2004) Alpha-7 nicotinic receptor agonists: potential new candidates for the treatment of schizophrenia. *Psychopharmacology (Berl)* 174:54-64.
- Marutle A, Zhang X, Court J, Piggott M, Johnson M, Perry R, Perry E, Nordberg A (2001) Laminar distribution of nicotinic receptor subtypes in cortical regions in schizophrenia. *J Chem Neuroanat* 22:115-126.
- Matteucci C (1842) Sur un phénomène physiologique produit par les muscles en contraction. *Ann chim et phys* 6:339.
- McGehee DS, Heath MJ, Gelber S, Devay P, Role LW (1995) Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science* 269:1692-1696.
- Meier J, Vannier C, Serge A, Triller A, Choquet D (2001) Fast and reversible trapping of surface glycine receptors by gephyrin. *Nat Neurosci* 4:253-260.
- Millar NS (2008) RIC-3: a nicotinic acetylcholine receptor chaperone. *Br J Pharmacol* 153 Suppl 1:S177-183.
- Nachmansohn D (1955) The generation of bioelectric potentials. *Circ Res* 3:429-433.
- Nader MA, Morgan D, Gage HD, Nader SH, Calhoun TL, Buchheimer N, Ehrenkaufer R, Mach RH (2006) PET imaging of dopamine D2 receptors during chronic cocaine self-administration in monkeys. *Nat Neurosci* 9:1050-1056.
- Newpher TM, Ehlers MD (2008) Glutamate receptor dynamics in dendritic microdomains. *Neuron* 58:472-497.
- Noda M, Takahashi H, Tanabe T, Toyosato M, Furutani Y, Hirose T, Asai M, Inayama S, Miyata T, Numa S (1982) Primary structure of alpha-subunit precursor of Torpedo californica acetylcholine receptor deduced from cDNA sequence. *Nature* 299:793-797.
- Noda M, Takahashi H, Tanabe T, Toyosato M, Kikuyotani S, Furutani Y, Hirose T, Takashima H, Inayama S, Miyata T, Numa S (1983a) Structural homology of Torpedo californica acetylcholine receptor subunits. *Nature* 302:528-532.
- Noda M, Takahashi H, Tanabe T, Toyosato M, Kikuyotani S, Hirose T, Asai M, Takashima H, Inayama S, Miyata T, Numa S (1983b) Primary structures of beta- and delta-subunit precursors of Torpedo californica acetylcholine receptor deduced from cDNA sequences. *Nature* 301:251-255.
- Nordberg A (1999) PET studies and cholinergic therapy in Alzheimer's disease. *Rev Neurol (Paris)* 155 Suppl 4:S53-63.
- Nourry C, Grant SG, Borg JP (2003) PDZ domain proteins: plug and play! *Sci STKE* 2003:RE7.
- Nuechterlein KH, Dawson ME (1984) Information processing and attentional functioning in the developmental course of schizophrenic disorders. *Schizophr Bull* 10:160-203.
- Nugent FS, Kauer JA (2008) LTP of GABAergic synapses in the ventral tegmental area and beyond. *J Physiol* 586:1487-1493.
- O'Neill MJ, Murray TK, Lakics V, Visanji NP, Duty S (2002) The role of neuronal nicotinic acetylcholine receptors in acute and chronic neurodegeneration. *Curr Drug Targets CNS Neurol Disord* 1:399-411.
- Oddo S, LaFerla FM (2006) The role of nicotinic acetylcholine receptors in Alzheimer's disease. *J Physiol Paris* 99:172-179.
- Olsen RW, Meunier JC, Changeux JP (1972) Progress in the purification of the cholinergic receptor protein from *Electrophorus electricus* by affinity chromatography. *FEBS Lett* 28:96-100.
- Palmer AM (2002) Pharmacotherapy for Alzheimer's disease: progress and prospects. *Trends Pharmacol Sci* 23:426-433.
- Paradis S, Harrar DB, Lin Y, Koon AC, Hauser JL, Griffith EC, Zhu L, Brass LF, Chen C, Greenberg ME (2007) An RNAi-based approach identifies molecules required for glutamatergic and GABAergic synapse development. *Neuron* 53:217-232.
- Parker MJ, Zhao S, Bredt DS, Sanes JR, Feng G (2004) PSD93 regulates synaptic stability at neuronal cholinergic synapses. *J Neurosci* 24:378-388.
- Pelkmans L, Zerlin M (2005) Kinase-regulated quantal assemblies and kiss-and-run recycling of caveolae. *Nature* 436:128-133.

- Peng X, Katz M, Gerzanich V, Anand R, Lindstrom J (1994) Human alpha 7 acetylcholine receptor: cloning of the alpha 7 subunit from the SH-SY5Y cell line and determination of pharmacological properties of native receptors and functional alpha 7 homomers expressed in *Xenopus* oocytes. *Mol Pharmacol* 45:546-554.
- Perez JL, Khatri L, Chang C, Srivastava S, Osten P, Ziff EB (2001) PICK1 targets activated protein kinase Calpha to AMPA receptor clusters in spines of hippocampal neurons and reduces surface levels of the AMPA-type glutamate receptor subunit 2. *J Neurosci* 21:5417-5428.
- Perry E, Martin-Ruiz C, Lee M, Griffiths M, Johnson M, Piggott M, Haroutunian V, Buxbaum JD, Nasland J, Davis K, Gotti C, Clementi F, Tzartos S, Cohen O, Soreq H, Jaros E, Perry R, Ballard C, McKeith I, Court J (2000) Nicotinic receptor subtypes in human brain ageing, Alzheimer and Lewy body diseases. *Eur J Pharmacol* 393:215-222.
- Perry EK (1986) The cholinergic hypothesis--ten years on. *Br Med Bull* 42:63-69.
- Pettit DL, Shao Z, Yakel JL (2001) beta-Amyloid(1-42) peptide directly modulates nicotinic receptors in the rat hippocampal slice. *J Neurosci* 21:RC120.
- Picciotto MR, Zoli M (2002) Nicotinic receptors in aging and dementia. *J Neurobiol* 53:641-655.
- Plank C, Schillinger U, Scherer F, Bergemann C, Remy JS, Krotz F, Anton M, Lausier J, Rosenecker J (2003) The magnetofection method: using magnetic force to enhance gene delivery. *Biol Chem* 384:737-747.
- Poincheval-Fuhrman S, Sara SJ (1993) Chronic nicotine ingestion improves radial arm maze performance in rats. *Behav Pharmacol* 4:535-539.
- Pons S, Fattore L, Cossu G, Tolu S, Porcu E, McIntosh JM, Changeux JP, Maskos U, Fratta W (2008) Crucial role of alpha4 and alpha6 nicotinic acetylcholine receptor subunits from ventral tegmental area in systemic nicotine self-administration. *J Neurosci* 28:12318-12327.
- Quick MW, Lester RA (2002) Desensitization of neuronal nicotinic receptors. *J Neurobiol* 53:457-478.
- Racz B, Blanpied TA, Ehlers MD, Weinberg RJ (2004) Lateral organization of endocytic machinery in dendritic spines. *Nat Neurosci* 7:917-918.
- Radcliffe KA, Dani JA (1998) Nicotinic stimulation produces multiple forms of increased glutamatergic synaptic transmission. *J Neurosci* 18:7075-7083.
- Rangwala F, Drisdell RC, Rakhilin S, Ko E, Atluri P, Harkins AB, Fox AP, Salman SS, Green WN (1997) Neuronal alpha-bungarotoxin receptors differ structurally from other nicotinic acetylcholine receptors. *J Neurosci* 17:8201-8212.
- Renner M, Specht CG, Triller A (2008) Molecular dynamics of postsynaptic receptors and scaffold proteins. *Curr Opin Neurobiol* 18:532-540.
- Ripoll N, Bronnec M, Bourin M (2004) Nicotinic receptors and schizophrenia. *Curr Med Res Opin* 20:1057-1074.
- Roth AL, Berg DK (2003) Large clusters of alpha7-containing nicotinic acetylcholine receptors on chick spinal cord neurons. *J Comp Neurol* 465:195-204.
- Rozsa B, Katona G, Kaszas A, Szipocs R, Vizi ES (2008) Dendritic nicotinic receptors modulate backpropagating action potentials and long-term plasticity of interneurons. *Eur J Neurosci* 27:364-377.
- Rubinson DA, Dillon CP, Kwiatkowski AV, Sievers C, Yang L, Kopinja J, Rooney DL, Zhang M, Ihrig MM, McManus MT, Gertler FB, Scott ML, Van Parijs L (2003) A lentivirus-based system to functionally silence genes in primary mammalian cells, stem cells and transgenic mice by RNA interference. *Nat Genet* 33:401-406.
- Sadasivam G, Willmann R, Lin S, Erb-Vogtli S, Kong XC, Ruegg MA, Fuhrer C (2005) Src-family kinases stabilize the neuromuscular synapse in vivo via protein interactions, phosphorylation, and cytoskeletal linkage of acetylcholine receptors. *J Neurosci* 25:10479-10493.
- Sage D, Neumann FR, Hediger F, Gasser SM, Unser M (2005) Automatic tracking of individual fluorescence particles: application to the study of chromosome dynamics. *IEEE Trans Image Process* 14:1372-1383.
- Salas R, Main A, Gangitano D, De Biasi M (2007) Decreased withdrawal symptoms but normal tolerance to nicotine in mice null for the alpha7 nicotinic acetylcholine receptor subunit. *Neuropharmacology* 53:863-869.
- Sassoe-Pognetto M, Fritschy JM (2000) Mini-review: gephyrin, a major postsynaptic protein of GABAergic synapses. *Eur J Neurosci* 12:2205-2210.
- Saxton MJ, Jacobson K (1997) Single-particle tracking: applications to membrane dynamics. *Annu Rev Biophys Biomol Struct* 26:373-399.
- Sbalzarini IF, Koumoutsakos P (2005) Feature point tracking and trajectory analysis for video imaging in cell biology. *J Struct Biol* 151:182-195.
- Schmiedeberg O, Koppe R (1869) Das Muskarin. Das giftige Alkaloid des Fliegenpilzes. Vogel, Leipzig.
- Schoepfer R, Conroy WG, Whiting P, Gore M, Lindstrom J (1990) Brain alpha-bungarotoxin binding protein cDNAs and MAbs reveal subtypes of this branch of the ligand-gated ion channel gene superfamily. *Neuron* 5:35-48.
- Schrader N, Kim EY, Winking J, Paulukat J, Schindelin H, Schwarz G (2004) Biochemical characterization of the high affinity binding between the glycine receptor and gephyrin. *J Biol Chem* 279:18733-18741.
- Seguela P, Wadiche J, Dineley-Miller K, Dani JA, Patrick JW (1993) Molecular cloning, functional properties, and distribution of rat brain alpha 7: a nicotinic cation channel highly permeable to calcium. *J Neurosci* 13:596-604.
- Shimomura O, Johnson FH, Saiga Y (1962) Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusa, *Aequorea*. *J Cell Comp Physiol* 59:223-239.
- Shoop RD, Martone ME, Yamada N, Ellisman MH, Berg DK (1999) Neuronal acetylcholine receptors with alpha7 subunits are concentrated on somatic spines for synaptic signaling in embryonic chick ciliary ganglia. *J Neurosci* 19:692-704.
- Shoop RD, Esquenazi E, Yamada N, Ellisman MH, Berg DK (2002) Ultrastructure of a somatic spine mat for nicotinic signaling in neurons. *J Neurosci* 22:748-756.
- Smith DB, Johnson KS (1988) Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene* 67:31-40.
- Songyang Z, Fanning AS, Fu C, Xu J, Marfatia SM, Chishti AH, Crompton A, Chan AC, Anderson JM, Cantley LC (1997) Recognition of unique carboxyl-terminal motifs by distinct PDZ domains. *Science* 275:73-77.
- Staudinger J, Lu J, Olson EN (1997) Specific interaction of the PDZ domain protein PICK1 with the COOH terminus of protein kinase C-alpha. *J Biol Chem* 272:32019-32024.



- Staudinger J, Zhou J, Burgess R, Elledge SJ, Olson EN (1995) PICK1: a perinuclear binding protein and substrate for protein kinase C isolated by the yeast two-hybrid system. *J Cell Biol* 128:263-271.
- Steinberg JP, Takamiya K, Shen Y, Xia J, Rubio ME, Yu S, Jin W, Thomas GM, Linden DJ, Hugarir RL (2006) Targeted in vivo mutations of the AMPA receptor subunit GluR2 and its interacting protein PICK1 eliminate cerebellar long-term depression. *Neuron* 49:845-860.
- Strochlic L, Cartaud A, Cartaud J (2005) The synaptic muscle-specific kinase (MuSK) complex: new partners, new functions. *Bioessays* 27:1129-1135.
- Takai T, Noda M, Mishina M, Shimizu S, Furutani Y, Kayano T, Ikeda T, Kubo T, Takahashi H, Takahashi T, et al. (1985) Cloning, sequencing and expression of cDNA for a novel subunit of acetylcholine receptor from calf muscle. *Nature* 315:761-764.
- Takeya R, Takeshige K, Sumimoto H (2000) Interaction of the PDZ domain of human PICK1 with class I ADP-ribosylation factors. *Biochem Biophys Res Commun* 267:149-155.
- Tardin C, Cognet L, Bats C, Lounis B, Choquet D (2003) Direct imaging of lateral movements of AMPA receptors inside synapses. *Embo J* 22:4656-4665.
- Tasneem A, Iyer LM, Jakobsson E, Aravind L (2005) Identification of the prokaryotic ligand-gated ion channels and their implications for the mechanisms and origins of animal Cys-loop ion channels. *Genome Biol* 6:R4.
- Temburni MK, Rosenberg MM, Pathak N, McConnell R, Jacob MH (2004) Neuronal nicotinic synapse assembly requires the adenomatous polyposis coli tumor suppressor protein. *J Neurosci* 24:6776-6784.
- Terashima A, Cotton L, Dev KK, Meyer G, Zaman S, Duprat F, Henley JM, Collingridge GL, Isaac JT (2004) Regulation of synaptic strength and AMPA receptor subunit composition by PICK1. *J Neurosci* 24:5381-5390.
- Tonges L, Lingor P, Egle R, Dietz GP, Fahr A, Bahr M (2006) Stearilated octaarginine and artificial virus-like particles for transfection of siRNA into primary rat neurons. *Rna* 12:1431-1438.
- Torres GE, Yao WD, Mohn AR, Quan H, Kim KM, Levey AI, Staudinger J, Caron MG (2001) Functional interaction between monoamine plasma membrane transporters and the synaptic PDZ domain-containing protein PICK1. *Neuron* 30:121-134.
- Torres R, Firestein BL, Dong H, Staudinger J, Olson EN, Hugarir RL, Brecht DS, Gale NW, Yancopoulos GD (1998) PDZ proteins bind, cluster, and synaptically colocalize with Eph receptors and their ephrin ligands. *Neuron* 21:1453-1463.
- Triller A, Choquet D (2008) New concepts in synaptic biology derived from single-molecule imaging. *Neuron* 59:359-374.
- Tsen G, Williams B, Allaire P, Zhou YD, Ikononov O, Kondova I, Jacob MH (2000) Receptors with opposing functions are in postsynaptic microdomains under one presynaptic terminal. *Nat Neurosci* 3:126-132.
- Ullian EM, McIntosh JM, Sargent PB (1997) Rapid synaptic transmission in the avian ciliary ganglion is mediated by two distinct classes of nicotinic receptors. *J Neurosci* 17:7210-7219.
- Umbriaco D, Watkins KC, Descarries L, Cozzari C, Hartman BK (1994) Ultrastructural and morphometric features of the acetylcholine innervation in adult rat parietal cortex: an electron microscopic study in serial sections. *J Comp Neurol* 348:351-373.
- Unwin N (2005) Refined structure of the nicotinic acetylcholine receptor at 4 Å resolution. *J Mol Biol* 346:967-989.
- Van Kampen M, Selbach K, Schneider R, Schiegel E, Boess F, Schreiber R (2004) AR-R 17779 improves social recognition in rats by activation of nicotinic  $\alpha 7$  receptors. *Psychopharmacology (Berl)* 172:375-383.
- Vernallis AB, Conroy WG, Berg DK (1993) Neurons assemble acetylcholine receptors with as many as three kinds of subunits while maintaining subunit segregation among receptor subtypes. *Neuron* 10:451-464.
- Vogels TP, Rajan K, Abbott LF (2005) Neural network dynamics. *Annu Rev Neurosci* 28:357-376.
- Vulpain EFA (1866) *Leçons sur la physiologie générale et comparée du système nerveux faites au Muséum d'histoire naturelle*. Paris: G Baillière.
- Wang HY, Lee DH, D'Andrea MR, Peterson PA, Shank RP, Reitz AB (2000)  $\beta$ -Amyloid(1-42) binds to  $\alpha 7$  nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology. *J Biol Chem* 275:5626-5632.
- Wang J, Jing Z, Zhang L, Zhou G, Braun J, Yao Y, Wang ZZ (2003) Regulation of acetylcholine receptor clustering by the tumor suppressor APC. *Nat Neurosci* 6:1017-1018.
- Ward JM, Cockcroft VB, Lunt GG, Smillie FS, Wonnacott S (1990) Methyllaconitine: a selective probe for neuronal  $\alpha$ -bungarotoxin binding sites. *FEBS Lett* 270:45-48.
- Watanabe SY, Albsoul-Younes AM, Kawano T, Itoh H, Kaziro Y, Nakajima S, Nakajima Y (1999) Calcium phosphate-mediated transfection of primary cultured brain neurons using GFP expression as a marker: application for single neuron electrophysiology. *Neurosci Res* 33:71-78.
- Wiesner A, Fuhrer C (2006) Regulation of nicotinic acetylcholine receptors by tyrosine kinases in the peripheral and central nervous system: same players, different roles. *Cell Mol Life Sci* 63:2818-2828.
- Wightman RM, Haynes CL (2004) Synaptic vesicles really do kiss and run. *Nat Neurosci* 7:321-322.
- Williams BM, Temburni MK, Levey MS, Bertrand S, Bertrand D, Jacob MH (1998) The long internal loop of the  $\alpha 3$  subunit targets nAChRs to subdomains within individual synapses on neurons in vivo. *Nat Neurosci* 1:557-562.
- Williams ME, Burton B, Urrutia A, Shcherbatko A, Chavez-Noriega LE, Cohen CJ, Aiyar J (2005) Ric-3 promotes functional expression of the nicotinic acetylcholine receptor  $\alpha 7$  subunit in mammalian cells. *J Biol Chem* 280:1257-1263.
- Willmann R, Pun S, Stallmach L, Sadasivam G, Santos AF, Caroni P, Fuhrer C (2006) Cholesterol and lipid microdomains stabilize the postsynapse at the neuromuscular junction. *Embo J* 25:4050-4060.
- Witkin JM, Marek GJ, Johnson BG, Schoepp DD (2007) Metabotropic glutamate receptors in the control of mood disorders. *CNS Neurol Disord Drug Targets* 6:87-100.
- Wong HC, Bourdelas A, Krauss A, Lee HJ, Shao Y, Wu D, Mlodzik M, Shi DL, Zheng J (2003) Direct binding of the PDZ domain of Dishevelled to a conserved internal sequence in the C-terminal region of Frizzled. *Mol Cell* 12:1251-1260.
- Woodruff-Pak DS, Green JT, Coleman-Valencia C, Pak JT (2000) A nicotinic cholinergic agonist (GTS-21) and eyeblink classical conditioning: acquisition, retention, and relearning in older rabbits. *Exp Aging Res* 26:323-336.

- Woolf NJ (1991) Cholinergic systems in mammalian brain and spinal cord. *Prog Neurobiol* 37:475-524.
- Xia J, Zhang X, Staudinger J, Huganir RL (1999) Clustering of AMPA receptors by the synaptic PDZ domain-containing protein PICK1. *Neuron* 22:179-187.
- Xia Z, Dudek H, Miranti CK, Greenberg ME (1996) Calcium influx via the NMDA receptor induces immediate early gene transcription by a MAP kinase/ERK-dependent mechanism. *J Neurosci* 16:5425-5436.
- Xiao B, Tu JC, Petralia RS, Yuan JP, Doan A, Breder CD, Ruggiero A, Lanahan AA, Wenthold RJ, Worley PF (1998) Homer regulates the association of group 1 metabotropic glutamate receptors with multivalent complexes of homer-related, synaptic proteins. *Neuron* 21:707-716.
- Xu J, Xia J (2006) Structure and function of PICK1. *Neurosignals* 15:190-201.
- Young JW, Crawford N, Kelly JS, Kerr LE, Marston HM, Spratt C, Finlayson K, Sharkey J (2007) Impaired attention is central to the cognitive deficits observed in alpha 7 deficient mice. *Eur Neuropsychopharmacol* 17:145-155.
- Zago WM, Massey KA, Berg DK (2006) Nicotinic activity stabilizes convergence of nicotinic and GABAergic synapses on filopodia of hippocampal interneurons. *Mol Cell Neurosci* 31:549-559.
- Zarei MM, Radcliffe KA, Chen D, Patrick JW, Dani JA (1999) Distributions of nicotinic acetylcholine receptor alpha7 and beta2 subunits on cultured hippocampal neurons. *Neuroscience* 88:755-764.
- Zeitelhofer M, Vessey JP, Xie Y, Tubing F, Thomas S, Kiebler M, Dahm R (2007) High-efficiency transfection of mammalian neurons via nucleofection. *Nat Protoc* 2:1692-1704.
- Zhang J, Berg DK (2007) Reversible inhibition of GABAA receptors by alpha7-containing nicotinic receptors on the vertebrate postsynaptic neurons. *J Physiol* 579:753-763.
- Zhang ZW, Coggan JS, Berg DK (1996) Synaptic currents generated by neuronal acetylcholine receptors sensitive to alpha-bungarotoxin. *Neuron* 17:1231-1240.
- Zhou FM, Wilson CJ, Dani JA (2002) Cholinergic interneuron characteristics and nicotinic properties in the striatum. *J Neurobiol* 53:590-605.

## Curriculum Vitae

Surname	Bürli
Name	Thomas
Address	Winterthurerstrasse 84 CH-8006 Zürich
Citizenship, Hometown	Swiss, Klingnau (AG)
Marital Status	single
Birth Date	24.02.1979

### Education

#### Studies

diploma in biochemistry ETH Zürich	Oct 1999 – Mar 2004
Dipl. Sci. ETH Zürich	

#### Schools

grammar school Kriegacker, MuttENZ (natural science program)	1995 - Dec 1998
middle school Engerfeld, Rheinfelden	1991 - 1995
primary school Magden	1986 - 1991

#### Courses

Basic Management Skills	Feb 2008
ZNZ Neuroscience Advanced Course II	2007
ZNZ Neuroscience Advanced Course I	2007
ZNZ Statistics Course	2006
ZNZ Neuroscience Introductory Course II	2006
ZNZ Neuroscience Introductory Course I	2005

### Army

basic military service, Fribourg	Feb - Mai 1999
----------------------------------	----------------

### Employment Record

BMA AG Augst, temporary employment as biochemist	May - Oct 2004
Feldschlösschen AG Rheinfelden, research assistant	Jun - Sep 2002
Rohner AG Pratteln, assistant	Jun - Sep 1999

## List of Publications

Single particle tracking of  $\alpha 7$  nicotinic acetylcholine receptors in hippocampal neurons reveals regulated perisynaptic confinement at glutamatergic and GABAergic perisynaptic sites, *Journal of Neuroscience* 2009, submitted, **T. Bürli et al.**

Postsynaptic mechanisms influence the formation of GABAergic synapses in hippocampal cultured neurons, *Neuroscience* 2009, submitted, *C. Petitjean, T. Bürli, C. Sidler, J.-M. Fritschy*

Efficient transfection of DNA or shRNA vectors into neurons using magnetofection. *Nat Protoc.* 2007;2(12):3090-101. **T. Buerli & C. Pellegrino et al.**

PICK1 interacts with alpha 7 neuronal nicotinic acetylcholine receptors and controls their clustering. *Mol Cell Neurosci.* 2007 Jun;35(2):339-55. Epub 2007 Mar 24. *K. Baer & T. Bürli et al.*

Caveolin-Stabilized Membrane Domains as Multifunctional Transport and Sorting Devices in Endocytic Membrane Traffic. *Cell* 2004, Volume 118, Issue 6, Pages 767-780 *L. Pelkmans, T. Bürli, M. Zerial, A. Helenius*

Analysis of interaction between caveolae-mediated endocytosis and endocytosis via clathrin coated pits. Diploma work 2003, **T.Bürli**, supervised by Dr. A. Tagawa , Lab of Prof. Dr. A. Helenius, ETH Zürich

The effect of active and dominant negative mutants of RabGTPases on the entry of Simian Virus 40. Semester work 2002, **T.Bürli**, supervised by Prof. Dr. L. Pelkmans, Lab of Prof. Dr. A. Helenius, ETH Zürich

### Presentations

Federation of European Neuroscience Societies FENS 2008 (Genf)	Dynamics of alpha 7 neuronal nicotinic acetylcholine receptors.
Society for Neuroscience SfN 2007 (San Diego, USA)	Dynamics of alpha 7 neuronal nicotinic acetylcholine receptors.
Neuroscience Center Zürich 2007 (Zürich):	Dynamics of alpha 7 neuronal nicotinic acetylcholine receptors.
Swiss Society for Neuroscience SSN 2007 (Bern):	PICK1 interacts with alpha 7 neuronal nicotinic acetylcholine receptors and controls their clustering.
Swiss Society for Neuroscience SSN 2006 (Basel):	PICK1 interacts with alpha 7 neuronal nicotinic acetylcholine receptors and controls their clustering.
Neuroscience Center Zürich 2006 (Zürich):	PICK1 interacts with alpha 7 neuronal nicotinic acetylcholine receptors and controls their clustering.
Neuroscience Center Zürich 2005 (Zürich):	PICK1 interacts with alpha 7 neuronal nicotinic acetylcholine receptors and controls their clustering.